

7

ANTIBODY COMPOSITION-PRODUCING CELL

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a cell for the production of an antibody molecule such as an antibody useful for various diseases, a fragment of the antibody and a fusion protein having the Fc region of the antibody or the like, a method for producing an antibody composition using the cell, the antibody composition and use thereof.

2. Brief description of the Background Art

Since antibodies have high binding activity, binding specificity and high stability in blood, their applications to diagnosis, prevention and treatment of various human diseases have been attempted [*Monoclonal Antibodies: Principles and Applications*, Wiley-Liss, Inc., Chapter 2.1 (1995)]. Also, production of a humanized antibody such as a human chimeric antibody or a human complementarity determining region (hereinafter referred to as "CDR")-grafted antibody from an antibody derived from an animal other than human have been attempted by using genetic recombination techniques. The human chimeric antibody is an antibody in which its antibody variable region (hereinafter referred to as "V region") is an antibody derived from an animal other than human and its constant region (hereinafter referred to as "C region") is derived from a human antibody. The human CDR-grafted antibody is an antibody in which the CDR of a human antibody is replaced by CDR of an antibody derived from an animal other than human.

It has been revealed that five classes, namely IgM, IgD, IgG, IgA and IgE, are present in antibodies derived from mammals. Antibodies of human IgG class are mainly used for the diagnosis, prevention and treatment of various

human diseases because they have functional characteristics such as long half-life in blood, various effector functions and the like [Monoclonal Antibodies: Principles and Applications, Wiley-Liss, Inc., Chapter 1 (1995)]. The human IgG class antibody is further classified into the following 4 subclasses: IgG1, IgG2, IgG3 and IgG4. A large number of studies have so far been conducted for antibody-dependent cell-mediated cytotoxic activity (hereinafter referred to as "ADCC activity") and complement-dependent cytotoxic activity (hereinafter referred to as "CDC activity") as effector functions of the IgG class antibody, and it has been reported that among antibodies of the human IgG class, the IgG1 subclass has the highest ADCC activity and CDC activity [Chemical Immunology, 65, 88 (1997)]. In view of the above, most of the anti-tumor humanized antibodies, including commercially available Rituxan and Herceptin, which require high effector functions for the expression of their effects, are antibodies of the human IgG1 subclass.

Expression of ADCC activity and CDC activity of the human IgG1 subclass antibodies requires binding of the Fc region of the antibody to an antibody receptor existing on the surface of an effector cell, such as a killer cell, a natural killer cell, an activated macrophage or the like (hereinafter referred to as "FcγR") and various complement components are bound. Regarding the binding, it has been suggested that several amino acid residues in the hinge region and the second domain of C region (hereinafter referred to as "Cy2 domain") of the antibody are important [Eur. J. Immunol., 23, 1098 (1993), Immunology, 86, 319 (1995), Chemical Immunology, 65, 88 (1997)] and that a sugar chain binding to the Cy2 domain [Chemical Immunology, 65, 88 (1997)] is also important.

Regarding the sugar chain, Boyd et al. have examined effects of a sugar chain on the ADCC activity and

CDC activity by treating a human CDR-grafted antibody CAMPATH-1H (human IgG1 subclass) produced by a Chinese hamster ovary cell (CHO cell) or a mouse myeloma NSO cell (NSO cell) with various sugar hydrolyzing enzymes, and reported that elimination of the non-reducing end sialic acid did not have influence upon both activities, but the CDC activity alone was affected by further removal of galactose residue and about 50% of the activity was decreased, and that complete removal of the sugar chain caused disappearance of both activities [*Molecular Immunol.*, 32, 1311 (1995)]. Also, Lifely et al. have analyzed the sugar chain bound to a human CDR-grafted antibody CAMPATH-1H (human IgG1 subclass) which was produced by CHO cell, NSO cell or rat myeloma YO cell, measured its ADCC activity, and reported that the CAMPATH-1H derived from YO cell showed the highest ADCC activity, suggesting that N-acetylglucosamine (hereinafter referred also to as "GlcNAc") at the bisecting position is important for the activity [*Glycobiology*, 5, 813 (1995); WO 99/54342]. These reports indicate that the structure of the sugar chain plays an important role in the effector functions of human antibodies of IgG1 subclass and that it is possible to prepare an antibody having more higher effector function by changing the structure of the sugar chain. However, actually, structures of sugar chains are various and complex, and it cannot be said that an actual important structure for the effector function was identified.

Sugar chains of glycoproteins are roughly divided into two types, namely a sugar chain which binds to asparagine (N-glycoside-linked sugar chain) and a sugar chain which binds to other amino acid such as serine, threonine (O-glycoside-linked sugar chain), based on the binding form to the protein moiety. The N-glycoside-linked sugar chains have various structures [*Biochemical Experimentation Method* 23 - Method for Studying

$\text{Man } \alpha 1 \rightarrow \begin{matrix} 6 \\ 3 \end{matrix} \text{Man } \beta 1 \rightarrow 4\text{GlcNAc } \beta 1 \rightarrow 4\text{GlcNAc} \quad (\text{I})$

In the Fc region of an antibody of an IgG type, two N-glycoside-linked sugar chain binding sites are present. In serum IgG, to the sugar chain binding site, generally, binds a complex type sugar chain having plural branches and in which addition of sialic acid or bisecting N-acetylglucosamine is low. It is known that there is variety regarding the addition of galactose to the non-reducing end of the complex type sugar chain and the addition of fucose to the N-acetylglucosamine in the reducing end [*Biochemistry*, 36, 130 (1997)].

It has been considered that such a structure of a sugar chain is determined by sugar chain genes, namely a gene for a glycosyltransferase which synthesizes a sugar chain and a gene for a glycolytic enzyme which hydrolyzes the sugar chain.

Synthesis of an *N*-glycoside-linked sugar chain is described below.

Glycoproteins are modified with a sugar chain in the endoplasmic reticulum (hereinafter referred to as "ER") lumen. During the biosynthesis step of the *N*-glycoside-linked sugar chain, a relatively large sugar chain is transferred to the polypeptide chain which is elongating in the ER lumen. In the transformation, the sugar chain is firstly added in succession to phosphate groups of a long chain lipid carrier comprising about 20 α -isoprene units, which is called dolichol phosphate (hereinafter referred also to as "P-Dol"). That is, *N*-acetylglucosamine is transferred to dolichol phosphate to thereby form GlcNAc-P-P-Dol and then one more GlcNAc is transferred to form GlcNAc-GlcNAc-P-P-Dol. Next, five mannoses (hereinafter mannose is also referred to as "Man") are transferred to thereby form (Man)₅-(GlcNAc)₂-P-P-Dol and then four Man's and three glucoses (hereinafter glucose is also referred to as "Glc") are transferred. Thus, a sugar chain precursor, (Glc)₃-(Man)₄-(GlcNAc)₂-P-P-Dol, called core oligosaccharide is formed. The sugar chain precursor comprising 14 sugars is transferred as a mass to a polypeptide having an asparagine-X-serine or asparagine-X-threonine sequence in the ER lumen. In the reaction, dolichol pyrophosphate (P-P-Dol) bound to the core oligosaccharide is released but again becomes dolichol phosphate by hydrolysis with pyrophosphatase and is recycled. Trimming of the sugar chain immediately starts after the sugar chain binds to the polypeptide. That is, 3 Glc's and 1 or 2 Man's are eliminated on the ER, and it is known that α -1,2-

glucosidase I, α -1,3-glucosidase II and α -1,2-mannosidase relates to the elimination.

The glycoprotein which was subjected to trimming on the ER is transferred to the Golgi body and are variously modified. In the cis part of the Golgi body, N-acetylglucosamine phosphotransferase which relates to addition of mannose phosphate, N-acetylglucosamine 1-phosphodiester α -N-acetylglucosaminidase and α -mannosidase I are present and reduce the Man residues to 5. In the medium part of the Golgi body, N-acetylglucosamine transferase I (GnTI) which relates to addition of the first outside GlcNAc of the complex type N-glycoside-linked sugar chain, α -mannosidase II which relates to elimination of 2 Man's, N-acetylglucosamine transferase II (GnTII) which relates to addition of the second GlcNAc from the outside and α -1,6-fucosyltransferase which relates to addition of fucose to the reducing end N-acetylglucosamine are present. In the trans part of the Golgi body, galactose transferase which relates to addition of galactose and sialyltransferase which relates to addition of sialic acid such as N-acetylneuraminic acid or the like are present. It is known that N-glycoside-linked sugar chain is formed by activities of these various enzymes.

In general, most of the humanized antibodies of which application to medicaments is in consideration are prepared using genetic recombination techniques and produced using Chinese hamster ovary tissue-derived CHO cell as the host cell. But as described above, since the sugar chain structure plays a remarkably important role in the effector function of antibodies and differences are observed in the sugar chain structure of glycoproteins expressed by host cells, development of a host cell which can be used for the production of an antibody having higher effector function is desired.

structures have been obtained as a lectin-resistant cell line using a lectin such as WGA (wheat-germ agglutinin derived from *T. vulgaris*), ConA (concanavalin A derived from *C. ensiformis*), RIC (a toxin derived from *R. communis*), L-PHA (leucoagglutinin derived from *P. vulgaris*), LCA (lentil agglutinin derived from *L. culinaris*), PSA (pea lectin derived from *P. sativum*) or the like [Somatic Cell Mol. Genet., 12, 51 (1986)].

As an example of the modification of the sugar chain structure of a product obtained by introducing the gene of an enzyme relating to the modification of a sugar chain into a host cell, it has been reported that a protein in which a number of sialic acid is added to the non-reducing end of the sugar chain can be produced by introducing rat β -galactoside- α -2,6-sialyltransferase into CHO cell [J. Biol. Chem., 261, 13848 (1989)].

Also, it was confirmed that an H antigen (Fucal-2Gal β 1-) in which fucose (hereinafter also referred to as "Fuc") was added to the non-reducing end of the sugar chain was expressed by introducing human β -galactoside-2- α -fucosyltransferase into mouse L cell [Science, 252, 1668 (1991)]. In addition, based on knowledge that addition of the bisecting-positioned N-acetylglucosamine of N-glycoside-linked sugar chain is important for the ADCC activity of antibody, Umana et al. have prepared CHO cell which expresses β -1,4-N-acetylglucosamine transferase III (GnTIII) and compared it with the parent cell line on the expression of GnTIII. It was confirmed that expression of GnTIII was not observed in the parent cell line of CHO cell [J. Biol. Chem., 261, 13370 (1984)], and that the antibody expressed using the produced GnTIII expressing CHO cell had ADCC activity 16 times higher than the antibody expressed using the parent cell line [Glycobiology, 5, 813 (1995): WO 99/54342]. At this time, Umana et al. have also produced CHO cell into which β -1,4-N-acetylglucosamine transferase V

(GnTV) was introduced and reported that excess expression of GnTIII or GnTV shows toxicity for CHO cell.

Thus, in order to modify the sugar chain structure of the glycoprotein to be produced, control of the activity of the enzyme relating to the modification of a sugar chain in the host cell has been attempted, but actually, the structures of sugar chains are various and complex, and solution of the physiological roles of sugar chains would be insufficient, so that trial and error are repeated. Particularly, although it has been revealed little by little that the effector function of antibodies is greatly influenced by the sugar chain structure, a truly important sugar chain structure has not been specified yet. Accordingly, identification of a sugar chain which has influence upon the effector function of antibodies and development of a host cell to which such a sugar chain structure can be added are expected for developing medicaments.

SUMMARY OF THE INVENTION

An object of the present invention is to provide a host cell which produces an antibody composition and can control a sugar chain structure bound to an antibody molecule, a cell which can produce an antibody composition having high ADCC activity, a production method of an antibody composition using the cell and an antibody composition produced by the production method.

The present invention relates to the following (1) to (61).

(1) A Chinese hamster ovary tissue-derived CHO cell into which a gene encoding an antibody molecule is introduced, which produces an antibody composition comprising an antibody molecule having complex N-glycoside-linked sugar chains bound to the Fc region, wherein among the total complex N-glycoside-linked sugar chains bound to

the Fc region in the composition, the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain is 20% or more.

(2) The CHO cell according to (1), wherein the sugar chain to which fucose is not bound is a complex *N*-glycoside-linked sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond.

(3) The CHO cell according to (1) or (2), wherein the antibody molecule belongs to an IgG class.

(4) The CHO cell according to any one of (1) to (3), wherein the activity of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose and/or the activity of an enzyme relating to the modification of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain is decreased or deleted.

(5) The CHO cell according to (4), wherein the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose is an enzyme selected from the group consisting of the following (a), (b) and (c):

(a) GMD (GDP-mannose 4,6-dehydratase);

(b) Fx (GDP-keto-6-deoxymannose 3,5-epimerase, 4-reductase);

(c) GFPP (GDP-beta-L-fucose pyrophosphorylase).

(6) The CHO cell according to (5), wherein the GMD is a protein encoded by a DNA of the following (a) or (b):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:65;

(b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:65 under stringent conditions and encodes a protein having GMD activity.

(7) The CHO cell according to (5), wherein the GMD is a protein selected from the group consisting of the following (a), (b) and (c):

(a) a protein comprising the amino acid sequence represented by SEQ ID NO:71;

(b) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:71 and has GMD activity;

(c) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:71 and has GMD activity.

(8) The CHO cell according to (5), wherein the Fx is a protein encoded by a DNA of the following (a) or (b):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:48;

(b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:48 under stringent conditions and encodes a protein having Fx activity.

(9) The CHO cell according to (5), wherein the Fx is a protein selected from the group consisting of the following (a), (b) and (c):

(a) a protein comprising the amino acid sequence represented by SEQ ID NO:72;

(b) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:72 and has Fx activity;

(c) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:72 and has Fx activity.

stringent conditions and encodes a protein having α -1,6-fucosyltransferase activity.

(14) The CHO cell according to (12), wherein the α -1,6-fucosyltransferase is a protein selected from the group consisting of the following (a), (b) and (c):

(a) a protein comprising the amino acid sequence represented by SEQ ID NO:23;

(b) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:23 and has α -1,6-fucosyltransferase activity;

(c) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:23 and has α -1,6-fucosyltransferase activity.

(15) The CHO cell according to any one of (4) to (14), wherein the enzyme activity is decreased or deleted by a technique selected from the group consisting of the following (a), (b), (c), (d) and (e):

(a) a gene disruption technique targeting a gene encoding the enzyme;

(b) a technique for introducing a dominant negative mutant of a gene encoding the enzyme;

(c) a technique for introducing mutation into the enzyme;

(d) a technique for inhibiting transcription or translation of a gene encoding the enzyme;

(e) a technique for selecting a cell line resistant to a lectin which recognizes a sugar chain in which 1-position of fucose is bound to 6-position of N-acetylglucosamine in the reducing end through α -bond in the complex N-glycoside-linked sugar chain.

(16) The CHO cell according to any one of (4) to (15), which is resistant to at least a lectin which recognizes a

(c) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:71 and has GMD activity.

(27) The cell according to (24), wherein the Fx is a protein encoded by a DNA of the following (a) or (b):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:48;

(b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:48 under stringent conditions and encodes a protein having Fx activity.

(28) The cell according to (24), wherein the Fx is a protein selected from the group consisting of the following (a), (b) and (c):

(a) a protein comprising the amino acid sequence represented by SEQ ID NO:72;

(b) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:72 and has Fx activity;

(c) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:72 and has Fx activity.

(29) The cell according to (24), wherein the GFPP is a protein encoded by a DNA of the following (a) or (b):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:51;

(b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:51 under stringent conditions and encodes a protein having GFPP activity.

(30) The cell according to (24), wherein the GFPP is a protein selected from the group consisting of the following (a), (b) and (c):

(a) a protein comprising the amino acid sequence represented by SEQ ID NO:73;

(b) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:73 and has GFPP activity;

(c) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:73 and has GFP activity.

(31) The cell according to (23), wherein the enzyme relating to the modification of a sugar chain in which 1-position of fucose is bound to 6-position of N-acetylglucosamine in the reducing end through α -bond in the N-glycoside-linked sugar chain is α -1,6-fucosyltransferase.

(32) The cell according to (31), wherein the α -1,6-fucosyltransferase is a protein encoded by a DNA selected from the group consisting of the following (a), (b), (c) and (d):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:1;

(b) a DNA comprising the nucleotide sequence represented by SEQ ID NO:2;

(c) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:1 under stringent conditions and encodes a protein having α -1,6-fucosyltransferase activity,;

(d) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:2 under stringent conditions and encodes a protein having α -1,6-fucosyltransferase activity.

(33) The cell according to (31), wherein the α -1,6-fucosyltransferase is a protein selected from the group consisting of the following (a), (b), (c), (d), (e) and (f):

(a) a protein comprising the amino acid sequence represented by SEQ ID NO:23;

(b) a protein comprising the amino acid sequence represented by SEQ ID NO:24;

(c) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:23 and has α -1,6-fucosyltransferase activity;

(d) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:24 and has α -1,6-fucosyltransferase activity;

(e) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:23 and has α -1,6-fucosyltransferase activity;

(f) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:24 and has α -1,6-fucosyltransferase activity.

(34) The cell according to any one of (23) to (33), wherein the genetic engineering technique is a technique selected from the group consisting of the following (a), (b), (c) and (d):

(a) a gene disruption technique targeting a gene encoding the enzyme;

(b) a technique for introducing a dominant negative mutant of a gene encoding the enzyme;

(c) a technique for introducing mutation into the enzyme;

(d) a technique for inhibiting transcription and/or translation of a gene encoding the enzyme.

(35) The cell according to any one of (23) to (34), which is resistant to at least a lectin which recognizes a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain.

(36) The cell according to any one of (23) to (35), which is a cell selected from the group consisting of the following (a) to (i):

(a) a CHO cell derived from a Chinese hamster ovary tissue;

(b) a rat myeloma cell line, YB2/3HL.P2.G11.16Ag.20 cell;

(c) a mouse myeloma cell line, NSO cell;

(d) a mouse myeloma cell line, SP2/0-Ag14 cell;

(e) a BHK cell derived from a syrian hamster kidney tissue;

(f) an antibody-producing hybridoma cell;

(g) a human leukemia cell line, Namalwa cell;

(h) an embryonic stem cell;

(i) a fertilized egg cell.

(37) The cell according to any one of (23) to (36) into which a gene encoding an antibody molecule is introduced.

(38) The cell according to (37), wherein the antibody molecule belongs to an IgG class.

(39) A method for producing an antibody composition, which comprises culturing the cell according to (37) or (38) in a medium to produce and accumulate the antibody composition in the culture; and recovering the antibody composition from the culture.

(40) The method according to (39), which produces an antibody composition having higher antibody-dependent cell-

mediated cytotoxic activity than an antibody composition obtained from its parent cell line.

(41) An antibody composition which is produced using the method according to (39) or (40).

(42) A transgenic non-human animal or plant or the progenies thereof, comprising a genome which is modified such that the activity of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose and/or the activity of an enzyme relating to the modification of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain is decreased.

(43) The transgenic non-human animal or plant or the progenies thereof according to (42), wherein a gene encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or a gene encoding the enzyme relating to the modification of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain is knocked out.

(44) The transgenic non-human animal or plant or the progenies thereof according to (42) or (43), wherein the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose is an enzyme selected from the group consisting of the following (a), (b) and (c):

(a) GMD (GDP-mannose 4,6-dehydratase);

(b) Fx (GDP-keto-6-deoxymannose 3,5-epimerase, 4-reductase);

(c) GFPP (GDP-beta-L-fucose pyrophosphorylase).

(45) The transgenic non-human animal or plant or the progenies thereof according to (44), wherein the GMD is a protein encoded by a DNA of the following (a) or (b):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:65;

(b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:65 under stringent conditions and encodes a protein having GMD activity.

(46) The transgenic non-human animal or plant or the progenies thereof according to (44), wherein the Fx is a protein encoded by a DNA of the following (a) or (b):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:48;

(b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:48 under stringent conditions and encodes a protein having Fx activity.

(47) The transgenic non-human animal or plant or the progenies thereof according to (44), wherein the GFPP is a protein encoded by a DNA of the following (a) or (b):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:51;

(b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:51 under stringent conditions and encodes a protein having GFPP activity.

(48) The transgenic non-human animal or plant or the progenies thereof according to (42) or (43), wherein the enzyme relating to the modification of a sugar chain in which 1-position of fucose is bound to 6-position of N-acetylglucosamine in the reducing end through α -bond in the N-glycoside-linked sugar chain is α -1,6-fucosyltransferase.

(49) The transgenic non-human animal or plant or the progenies thereof according to (48), wherein the α -1,6-fucosyltransferase is a protein encoded by a DNA selected from the group consisting of the following (a), (b), (c) and (d):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:1;

(b) a DNA comprising the nucleotide sequence represented by SEQ ID NO:2;

(c) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:1 under stringent conditions and encodes a protein having α -1,6-fucosyltransferase activity;

(d) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:2 under stringent conditions and encodes a protein having α -1,6-fucosyltransferase activity.

(50) The transgenic non-human animal or plant or the progenies thereof according to any one of (42) to (49), wherein the transgenic non-human animal is an animal selected from the group consisting of cattle, sheep, goat, pig, horse, mouse, rat, fowl, monkey and rabbit.

(51) A method for producing an antibody composition, which comprises introducing a gene encoding an antibody molecule into the transgenic non-human animal or plant or the progenies thereof according to any one of (42) to (50); rearing the animal or plant; isolating tissue or body fluid comprising the introduced antibody from the reared animal or plant; and recovering the antibody composition from the isolated tissue or body fluid.

(52) The method according to (51), wherein the antibody molecule belongs to an IgG class.

(53) The method according to (51) or (52), which produces an antibody composition having higher antibody-dependent cell-mediated cytotoxic activity than an antibody composition obtained from a non-human animal or plant or the progenies thereof whose genome is not modified.

(54) An antibody composition which is produced using the method according to any one of (51) to (53).

(55) A medicament comprising the antibody composition according to any one of (21), (22), (41) and (54) as an active ingredient.

(56) The medicament according to (55), wherein the medicament is a diagnostic drug, a preventive drug or a therapeutic drug for diseases accompanied by tumors, diseases accompanied by allergies, diseases accompanied by inflammations, autoimmune diseases, circulatory organ diseases, diseases accompanied by viral infections or diseases accompanied by bacterial infections.

(57) A protein selected from the group consisting of the following (a), (b), (c), (d), (e), (f), (g), (h), (i) and (j):

(a) a protein comprising the amino acid sequence represented by SEQ ID NO:71;

(b) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:71 and has GMD activity;

(c) a protein comprising the amino acid sequence represented by SEQ ID NO:72;

(d) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:72 and has Fx activity;

(e) a protein comprising the amino acid sequence represented by SEQ ID NO:73;

(f) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:73 and has GFPP activity;

(g) a protein comprising the amino acid sequence represented by SEQ ID NO:23;

(h) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:23 and has α -1,6-fucosyltransferase activity;

(i) a protein comprising the amino acid sequence represented by SEQ ID NO:24;

(j) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:24 and the α -1,6-fucosyltransferase activity.

(58) A DNA which encodes the protein according to (57).

(59) A DNA selected from the group consisting of the following (a), (b), (c), (d) and (e):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:1;

(b) a DNA comprising the nucleotide sequence represented by SEQ ID NO:2;

(c) a DNA comprising the nucleotide sequence represented by SEQ ID NO:65;

(d) a DNA comprising the nucleotide sequence represented by SEQ ID NO:48;

(e) a DNA comprising the nucleotide sequence represented by SEQ ID NO:51.

(60) A genome DNA selected from the group consisting of the following (a), (b) and (c):

(a) a genome DNA comprising the nucleotide sequence represented by SEQ ID NO:3;

(b) a genome DNA comprising the nucleotide sequence represented by SEQ ID NO:67;

(c) a genome DNA comprising the nucleotide sequence represented by SEQ ID NO:70.

(61) A target vector for homologous recombination, comprising a full length of the DNA according to any one of (58) to (60), or a part thereof.

BRIEF EXPLANATION OF THE DRAWINGS

Fig. 1 shows electrophoresis patterns of SDS-PAGE of five purified anti-GD3 chimeric antibodies (using

gradient gel from 4 to 15%). Fig. 1A and Fig. 1B show a result of the electrophoresis under non-reducing conditions and that under reducing conditions, respectively. Lanes 1 to 7 show electrophoresis patterns of high molecular weight markers, YB2/0-GD3 chimeric antibody, CHO/DG44-GD3 chimeric antibody, SP2/0-GD3 chimeric antibody, NS0-GD3 chimeric antibody (302), NS0-GD3 chimeric antibody (GIT) and low molecular weight markers, respectively.

Fig. 2 shows activities of five purified anti-GD3 chimeric antibodies to bind to GD3, measured by changing the antibody concentration. The ordinate and the abscissa show the binding activity with GD3 and the antibody concentration, respectively. "○", "●", "□", "■" and "△" show the activities of YB2/0-GD3 chimeric antibody, CHO/DG44-GD3 chimeric antibody, SP2/0-GD3 chimeric antibody, NS0-GD3 chimeric antibody (302) and NS0-GD3 chimeric antibody (GIT), respectively.

Fig. 3 shows ADCC activities of five purified anti-GD3 chimeric antibodies for a human melanoma cell line G-361. The ordinate and abscissa show the cytotoxic activity and the antibody concentration, respectively. "○", "●", "□", "■" and "△" show the activities of YB2/0-GD3 chimeric antibody, CHO/DG44-GD3 chimeric antibody, SP2/0-GD3 chimeric antibody, NS0-GD3 chimeric antibody (302) and NS0-GD3 chimeric antibody (GIT), respectively.

Fig. 4 shows electrophoresis patterns of SDS-PAGE of three purified anti-hIL-5Rα CDR-grafted antibodies (using gradient gel from 4 to 15%). Fig. 4A and Fig. 4B show results of the electrophoresis carried out under non-reducing conditions and those under reducing conditions, respectively. Lanes 1 to 5 show electrophoresis patterns of high molecular weight markers, YB2/0-hIL-5R CDR antibody, CHO/d-hIL-5R CDR antibody, NS0-hIL-5R CDR antibody and low molecular weight markers, respectively.

Fig. 5 shows activities of three purified anti-hIL-5R α CDR-grafted antibodies to bind to hIL-5R α , measured by changing the antibody concentration. The ordinate and the abscissa show the binding activity with hIL-5R α and the antibody concentration, respectively. "○", "●" and "□" show the activities of YB2/0-hIL-5R CDR antibody, CHO/d-hIL-5R CDR antibody and NS0-hIL-5R CDR antibody, respectively.

Fig. 6 show ADCC activities of three purified anti-hIL-5R α CDR-grafted antibodies for an hIL-5R expressing mouse T cell line CTLL-2(h5R). The ordinate and the abscissas show the cytotoxic activity and the antibody concentration, respectively. "○", "●" and "□" show the activities of YB2/0-hIL-5RCR antibody, CHO/d-hIL-5R CDR antibody and NS0-hIL-5R CDR antibody, respectively.

Fig. 7 shows inhibition activities of three purified anti-hIL-5R α CDR-grafted antibodies in an hIL-5-induced eosinophil increasing model of *Macaca fascicularis*. The ordinate and the abscissa show the number of eosinophils in peripheral blood and the number of days (the day of the commencement of antibody and hIL-5 administration was defined as 0 day). "101 and 102", "301, 302 and 303", "401, 402 and 403" and "501, 502 and 503" show results in the antibody non-administration group, the YB2/0-hIL-5R CDR antibody administered group, the CHO/d-hIL-5R CDR antibody administered group and the NS0-hIL-5R CDR antibody administered group, respectively.

Fig. 8 shows elution patterns of reverse phase HPLC elution of a PA-treated sugar chain (left side), and an elution pattern obtained by treating the PA-treated sugar chain with α -L-fucosidase and then analyzed by reverse phase HPLC (right side), of the purified anti-hIL-5R α CDR-grafted antibody produced by YB2/0 (Fig. 8A) and the purified anti-hIL-5R α CDR-grafted antibody produced by NS0 (Fig. 8B). The ordinates and the abscissas show the

relative fluorescence intensity and the elution time, respectively.

Fig. 9 shows an elution pattern obtained by preparing a PA-treated sugar chain from the purified anti-hIL-5R α CDR-grafted antibody produced by CHO/d cell and analyzing it by reverse phase HPLC. The ordinate and the abscissa show the relative fluorescence intensity and the elution time, respectively.

In Fig. 10, Fig. 10A shows the GD3-binding activities of a non-adsorbed fraction and a part of an adsorbed fraction, measured by changing the antibody concentration. The ordinate and the abscissa show the binding activity with GD3 and the antibody concentration, respectively. "●" and "○" show the non-adsorbed fraction and a part of the adsorbed fraction, respectively. Fig. 10B shows the ADCC activities of the non-adsorbed fraction and a part of the adsorbed fraction for a human melanoma line G-361. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively. "●" and "○" show the non-adsorbed fraction and a part of the adsorbed fraction, respectively.

Fig. 11 shows elution patterns obtained by analyzing PA-treated sugar chains prepared from a non-adsorbed fraction and a part of an adsorbed fraction by a reverse HPLC. Fig. 11A and Fig. 11B show an elution pattern of the non-adsorbed fraction and an elution pattern of a part of the adsorbed fraction, respectively. The ordinates and the abscissas show the relative fluorescence strength and the elution time, respectively.

Fig. 12 shows elution patterns of PA-treated sugar chains prepared from 6 anti-GD3 chimeric antibodies (Fig. 12A to Fig. 12F), obtained by analyzing them by reverse phase HPLC. The ordinates and the abscissas show the relative fluorescence intensity and the elution time, respectively.

Fig. 16 shows elution patterns of PA-treated sugar chains prepared from six kinds of anti-GD3 chimeric

antibodies, obtained by analyzing them by reverse phase HPLC. The ordinates and the abscissas show the relative fluorescence intensity and the elution time, respectively.

Fig. 17 shows CCR4-binding activities of six kinds of anti-CCR4 chimeric antibodies having a different ratio of α -1,6-fucose-free sugar chains measured by changing the antibody concentration. The ordinate and the abscissa show the binding activity with CCR4 and the antibody concentration, respectively. "■", "□", "▲", "△", "●" and "○" show the activities of anti-CCR4 chimeric antibody (46%), anti-CCR4 chimeric antibody (39%), anti-CCR4 chimeric antibody (27%), anti-CCR4 chimeric antibody (18%), anti-CCR4 chimeric antibody (9%) and anti-CCR4 chimeric antibody (8%), respectively.

Fig. 18 shows ADCC activities of anti-CCR4 chimeric antibodies having a different ratio of α -1,6-fucose-free sugar chains against CCR4/EL-4 cell, using an effector cell of the donor A. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively. "■", "□", "▲", "△", "●" and "○" show the activities of anti-CCR4 chimeric antibody (46%), anti-CCR4 chimeric antibody (39%), anti-CCR4 chimeric antibody (27%), anti-CCR4 chimeric antibody (18%), anti-CCR4 chimeric antibody (9%) and anti-CCR4 chimeric antibody (8%), respectively. Also, Fig. 18A and Fig. 18B show results obtained using effector cells of the donor A and the donor B, respectively.

Fig. 19 shows ADCC activities of anti-CCR4 chimeric antibodies having a different ratio of α -1,6-fucose-free sugar chain against CCR4/EL-4 cell, using an effector cell of the donor B. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively. "■", "□", "▲", "△", "●" and "○" show the activities of anti-CCR4 chimeric antibody (46%), anti-CCR4 chimeric antibody (39%), anti-CCR4 chimeric antibody (27%),

by reverse phase HPLC. Fig. 30A and Fig. 30B show elution patterns of PA-treated sugar chains prepared from an antibody produced by mfFUT8-6-introduced cell line and PA-treated sugar chains prepared from an antibody produced by PAGE249-introduced cell line, respectively. The ordinate and the abscissa show the relative fluorescence intensity and the elution time, respectively.

Fig. 31 shows an elution pattern of PA-treated sugar chains prepared from Herceptin, obtained by analyzing them by reverse phase HPLC. The ordinate and the abscissa show the relative fluorescence intensity and the elution time, respectively.

Fig. 32 shows construction of a plasmid CHfFUT8-pCR2.1.

Fig. 33 shows construction of a plasmid ploxPPuro.

Fig. 34 shows construction of a plasmid pKOFUT8gE2-1.

Fig. 35 shows construction of a plasmid pKOFUT8gE2-2.

Fig. 36 shows construction of a plasmid pscFUT8gE2-3.

Fig. 37 shows construction of a plasmid pKOFUT8gE2-3.

Fig. 38 shows construction of a plasmid pKOFUT8gE2-4.

Fig. 39 shows construction of a plasmid pKOFUT8gE2-5.

Fig. 40 shows construction of a plasmid pKOFUT8Puro.

Fig. 41 shows results of genome Southern analyses of 1st. Δ FUT8 2-46-1 and 1st. Δ FUT8 2-46 as α -1,6-fucosyltransferase gene-disrupted CHO cell lines.

Fig. 42 shows ADCC activities of an anti-CCR4 chimeric antibody purified from an FUT8 allele gene-disrupted cell line. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration. "▲"

and "■" show the activities of a purified antibody derived from an anti-CCR4 chimeric antibody-producing CHO cell 5-03 and a purified antibody derived from 1st.ΔFUT8 2-46-1, respectively.

Fig. 43 shows ADCC activities of anti-CCR4 human chimeric antibodies produced by lectin-resistant cell lines. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration. "□", "■", "◆" and "▲" show the activities of antibodies produced by the strain 5-03, CHO/CCR4-LCA, CHO/CCR4-AAL and CHO/CCR4-PHA, respectively.

Fig. 44 shows ADCC activities of anti-CCR4 human chimeric antibodies produced by lectin-resistant cell lines. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively. "□", "△" and "●" show activities of antibodies produced by YB2/0 (KM2760 # 58-35-16), 5-03 and CHO/CCR4-LCA, respectively.

Fig. 45 shows elution patterns of PA-treated sugar chains prepared from purified anti-CCR4 human chimeric antibodies, obtained by analyzing them by reverse phase HPLC. The ordinate and the abscissa show the relative fluorescence intensity and the elution time, respectively. Fig. 45A, Fig. 45B, Fig. 45C and Fig. 45D show results of analyses of an antibody produced by the strain 5-03, an antibody produced by CHO/CCR4-LCA, an antibody produced by CHO/CCR4-AAL and an antibody produced by CHO/CCR4-PHA, respectively.

Fig. 46 shows the 1st step of construction of an expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 47 shows the 2nd step of construction of the expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 48 shows the 3rd step of construction of the expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 49 shows the 4th step of construction of the expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 50 shows the 5th step of construction of the expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 51 shows the 6th step of construction of the expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 52 shows resistance of GMD-expressed CHO/CCR4-LCA for LCA lectin. The measurement was carried out twice by defining the survival rate of a group of cells cultured without adding LCA lectin as 100%. In the drawing, "249" shows the survival rate of the CHO/CCR4-LCA introduced with an expression vector pAGE249 for LCA lectin. GMD shows resistance of the CHO/CCR4-LCA introduced with a GMD expression vector pAGE249GMD for LCA lectin.

Fig. 53 shows ADCC activities of an anti-CCR4 chimeric antibody produced by cells of GMD-expressed CHO/CCR4-LCA cell lines. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively.

Fig. 54 show a production step of a plasmid CHO-GMD in which the 5'-terminal of a clone 34-2 is introduced into the 5'-terminal of a CHO cell-derived GMD cDNA clone 22-8.

Fig. 55 shows an elution pattern of PA-treated sugar chains prepared from an anti-CCR4 human chimeric antibody purified from GMD gene-expressed CHO/CCR4-LCA, obtained by analyzing them by reverse phase HPLC. The ordinate and the abscissa show the relative fluorescence intensity and the elution time, respectively.

DETAILED DESCRIPTION OF THE INVENTION

The Chinese hamster ovary tissue-derived CHO cell into which a gene encoding an antibody molecule is introduced according to the present invention may be any CHO cell, so long as it is a Chinese hamster ovary tissue-derived CHO cell into which a gene encoding an antibody molecule is introduced, which produces an antibody composition comprising complex N-glycoside-linked sugar chains bound to the Fc region of an antibody molecule, wherein among the total complex N-glycoside-linked sugar chains bound to the Fc region in the composition, the ratio of a sugar chain in which fucose is not bound to N-acetylglucosamine in the reducing end in the sugar chain is 20% or more.

In the present invention, the antibody molecule includes any molecule, so long as it comprises the Fc region of an antibody. Examples include an antibody, an antibody fragment, a fusion protein comprising an Fc region, and the like.

The antibody is a protein which is produced in the living body by immune reaction as a result of exogenous antigen stimulation and has an activity to specifically bind to the antigen. Examples of the antibody include an antibody secreted by a hybridoma cell prepared from a spleen cell of an animal immunized with an antigen; an antibody prepared by a genetic recombination technique, namely an antibody obtained by introducing an antibody gene-inserted antibody expression vector into a host cell; and the like. Specific examples include an antibody produced by a hybridoma, a humanized antibody, a human antibody and the like.

A hybridoma is a cell which is obtained by cell fusion between a B cell obtained by immunizing a mammal other than human with an antigen and a myeloma cell derived

from mouse or the like and can produce a monoclonal antibody having the desired antigen specificity.

Examples of the humanized antibody include a human chimeric antibody, a human CDR-grafted antibody and the like.

A human chimeric antibody is an antibody which comprises an antibody heavy chain variable region (hereinafter referred to as "HV" or "VH", the heavy chain being "H chain") and an antibody light chain variable region (hereinafter referred to as "LV" or "VL", the light chain being "L chain"), both of an animal other than human, a human antibody heavy chain constant region (hereinafter also referred to as "CH") and a human antibody light chain constant region (hereinafter also referred to as "CL"). As the animal other than human, any animal such as mouse, rat, hamster, rabbit or the like can be used, so long as a hybridoma can be prepared therefrom.

The human chimeric antibody can be produced by obtaining cDNA's encoding VH and VL from a monoclonal antibody-producing hybridoma, inserting them into an expression vector for host cell having genes encoding human antibody CH and human antibody CL to thereby construct a human chimeric antibody expression vector, and then introducing the vector into a host cell to express the antibody.

As the CH of human chimeric antibody, any CH can be used, so long as it belongs to human immunoglobulin (hereinafter referred to as "hIg") can be used. But those belonging to the hIgG class are preferable and any one of the subclasses belonging to the hIgG class, such as hIgG1, hIgG2, hIgG3 and hIgG4, can be used. Also, as the CL of human chimeric antibody, any CL can be used, so long as it belongs to the hIg class, and those belonging to the κ class or λ class can also be used.

A human CDR-grafted antibody is an antibody in which amino acid sequences of CDR's of VH and VL of an antibody derived from an animal other than human are grafted into appropriate positions of VH and VL of a human antibody.

The human CDR-grafted antibody can be produced by constructing cDNA's encoding V regions in which CDR's of VH and VL of an antibody derived from an animal other than human are grafted into CDR's of VH and VL of a human antibody, inserting them into an expression vector for host cell having genes encoding human antibody CH and human antibody CL to thereby construct a human CDR-grafted antibody expression vector, and then introducing the expression vector into a host cell to express the human CDR-grafted antibody.

As the CH of human CDR-grafted antibody, any CH can be used, so long as it belongs to the hIg, but those of the hIgG class are preferable and any one of the subclasses belonging to the hIgG class, such as hIgG1, hIgG2, hIgG3 and hIgG4, can be used. Also, as the CL of human CDR-grafted antibody, any CL can be used, so long as it belongs to the hIg class, and those belonging to the κ class or λ class can also be used.

A human antibody is originally an antibody naturally existing in the human body, but it also includes antibodies obtained from a human antibody phage library, a human antibody-producing transgenic animal and a human antibody-producing transgenic plant, which are prepared based on the recent advance in genetic engineering, cell engineering and developmental engineering techniques.

Regarding the antibody existing in the human body, a lymphocyte capable of producing the antibody can be cultured by isolating a human peripheral blood lymphocyte, immortalizing it by its infection with EB virus or the like

and then cloning it, and the antibody can be purified from the culture.

The human antibody phage library is a library in which antibody fragments such as Fab, single chain antibody and the like are expressed on the phage surface by inserting a gene encoding an antibody prepared from a human B cell into a phage gene. A phage expressing an antibody fragment having the desired antigen binding activity can be recovered from the library, using its activity to bind to an antigen-immobilized substrate as the marker. The antibody fragment can be converted further into a human antibody molecule comprising two full H chains and two full L chains by genetic engineering techniques.

A human antibody-producing transgenic non-human animal is an animal in which a human antibody gene is introduced into cells. Specifically, a human antibody-producing transgenic animal can be prepared by introducing a human antibody gene into ES cell of a mouse, transplanting the ES cell into an early stage embryo of other mouse and then developing it. By introducing a human chimeric antibody gene into a fertilized egg and developing it, the transgenic animal can be also prepared. Regarding the preparation method of a human antibody from the human antibody-producing transgenic animal, the human antibody can be produced and accumulated in a culture by obtaining a human antibody-producing hybridoma by a hybridoma preparation method usually carried out in mammals other than human and then culturing it.

Examples of the transgenic non-human animal include cattle, sheep, goat, pig, horse, mouse, rat, fowl, monkey, rabbit and the like.

Also, in the present invention, it is preferable that the antibody is an antibody which recognizes a tumor-related antigen, an antibody which recognizes an allergy- or inflammation-related antigen, an antibody which

recognizes circulatory organ disease-related antigen, an antibody which recognizes an autoimmune disease-related antigen or an antibody which recognizes a viral or bacterial infection-related antigen, and a human antibody which belongs to the IgG class is preferable.

An antibody fragment is a fragment which comprises the Fc region of an antibody. Examples of the antibody fragment include an H chain monomer, an H chain dimer and the like.

A fusion protein comprising an Fc region is a composition in which an antibody comprising the Fc region of an antibody or the antibody fragment is fused with a protein such as an enzyme, a cytokine or the like.

In the present invention, examples of the sugar chain which binds to the Fc region of an antibody molecule includes an *N*-glycoside-linked sugar chain. Examples of the *N*-glycoside-linked sugar chain include a complex type in which the non-reducing end side of the core structure has one or plural parallel branches of galactose-*N*-acetylglucosamine (hereinafter referred to as "Gal-GlcNAc") and the non-reducing end side of Gal-GlcNAc has a structure such as sialic acid, bisecting *N*-acetylglucosamine or the like.

In one antibody, the Fc region has positions to which an *N*-glycoside-linked sugar chain is bound which will be described later. Accordingly, two sugar chains are bound per one antibody molecule. Since the *N*-glycoside-linked sugar chain which binds to an antibody includes any sugar chain having the core structure represented by the structural formula (I), a number of combinations of sugar chains may possible for the two *N*-glycoside-linked sugar chains which bind to the antibody. Accordingly, identity of substances can be judged from the viewpoint of the sugar structure bound to the Fc region.

In the present invention, the composition which comprises an antibody molecule having complex *N*-glycoside-linked sugar chains in the Fc region (hereinafter referred to as "the antibody composition of the present invention") may comprise an antibody having the same sugar chain structure or an antibody having different sugar chain structures, so long as the effect of the present invention is obtained from the composition.

In the present invention, the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain among the total complex *N*-glycoside-linked sugar chains bound to the Fc region contained in the antibody composition is a ratio of the number of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain to the total number of the complex *N*-glycoside-linked sugar chains bound to the Fc region contained in the composition.

In the present invention, the sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is a sugar chain in which the fucose is not bound to *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain. Examples include a complex *N*-glycoside-linked sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine through α -bond.

The antibody composition shows high ADCC activity when the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain among the total complex *N*-glycoside-linked sugar chains binding to the Fc region contained in the antibody composition of the present invention is preferably 20% or more, more preferably 25% or more, still more preferably 30% or more, far preferably 40% or more, and most preferably 50% or more. As the antibody concentration

In the present invention, examples of the GMD include:

a protein encoded by a DNA of the following (a) or (b):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:65;

(b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:65 under stringent conditions and encodes a protein having GMD activity,

(c) a protein comprising the amino acid sequence represented by SEQ ID NO:71,

(d) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:71 and has GMD activity,

(e) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:71 and has GMD activity, and the like.

Also, examples of the DNA encoding the amino acid sequence of GMD include a DNA comprising the nucleotide sequence represented by SEQ ID NO:65 and a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:65 under stringent conditions and encodes an amino acid sequence having GMD activity.

In the present invention, examples of the Fx include:

a protein encoded by a DNA of the following (a) or (b):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:48;

(b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:48 under

stringent conditions and encodes a protein having Fx activity,

(c) a protein comprising the amino acid sequence represented by SEQ ID NO:72,

(d) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:72 and has Fx activity,

(e) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:72 and has Fx activity, and the like.

Also, examples of the DNA encoding the amino acid sequence of Fx include a DNA comprising the nucleotide sequence represented by SEQ ID NO:48 and a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:48 under stringent conditions and encodes an amino acid sequence having Fx activity.

In the present invention, examples of the GFPP include:

a protein encoded by a DNA of the following (a) or (b):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:51;

(b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:51 under stringent conditions and encodes a protein having GFPP activity,

(c) a protein comprising the amino acid sequence represented by SEQ ID NO:73,

(d) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:73 and has GFPP activity,

(e) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:73 and has GFPP activity, and the like.

Also, examples of the DNA encoding the amino acid sequence of GFPP include a DNA comprising the nucleotide sequence represented by SEQ ID NO:51 and a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:51 under stringent conditions and encodes an amino acid sequence having Fx activity.

In the present invention, the enzyme relating to the modification of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain includes any enzyme, so long as it is an enzyme relating to the reaction of binding of 1-position of fucose to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain.

Examples of the enzyme relating to the reaction of binding of 1-position of fucose to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain include α -1,6-fucosyltransferase, α -L-fucosidase and the like.

Also, examples include an enzyme which has influence on the activity the enzyme relating to the reaction of binding of 1-position of fucose to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain and an enzyme which has influence on the structure of substances as the substrate of the enzyme.

In the present invention, examples of the α -1,6-fucosyltransferase include:

a protein encoded by a DNA of the following (a), (b), (c) or (d):

(a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:1;

(b) a DNA comprising the nucleotide sequence represented by SEQ ID NO:2;

(c) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:1 under stringent conditions and encodes a protein having α -1,6-fucosyltransferase activity;

(d) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:2 under stringent conditions and encodes a protein having α -1,6-fucosyltransferase activity;

(e) a protein comprising the amino acid sequence represented by SEQ ID NO:23,

(f) a protein comprising the amino acid sequence represented by SEQ ID NO:24,

(g) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:23 and has α -1,6-fucosyltransferase activity,

(h) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:24 and has α -1,6-fucosyltransferase activity,

(i) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:23 and has α -1,6-fucosyltransferase activity,

(j) a protein which comprises an amino acid sequence having a homology of at least 80% with the amino acid sequence represented by SEQ ID NO:24 and has α -1,6-fucosyltransferase activity, and the like.

In the present invention, the protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:23, 24, 71, 72 or 73 and has α -1,6-fucosyltransferase activity, GMD activity, Fx activity or GFPP activity can be obtained, e.g., by introducing a site-directed mutation into a DNA encoding a protein having the amino acid sequence represented by SEQ ID NO:1, 2, 65, 48 or 51, respectively, using the site-directed mutagenesis described, e.g., in *Molecular Cloning*, Second Edition; *Current Protocols in Molecular Biology*; *Nucleic Acids Research*, 10, 6487 (1982); *Proc. Natl. Acad. Sci. USA*, 79, 6409 (1982); *Gene*, 34, 315 (1985); *Nucleic Acids Research*, 13, 4431 (1985); *Proc. Natl. Acad. Sci. USA*, 82, 488 (1985); and the like. The number of amino acids to be deleted, substituted, inserted and/or added is one or more, and the number is not particularly limited, but is a number which can be deleted, substituted or added by a known technique such as the site-directed mutagenesis, e.g., it is 1 to several tens, preferably 1 to 20, more preferably 1 to 10, and most preferably 1 to 5.

Also, in order to maintain the α -1,6-fucosyltransferase activity, GMD activity, Fx activity or GFPP activity of the protein to be used in the present invention, it has at least 80% or more, preferably 85% or more, more preferably 90% or more, still more preferably 95% or more, far more preferably 97% or more, and most preferably 99% or more, of homology with the amino acid sequence represented by SEQ ID NO:23, 24, 71, 72 or 73, when calculated using an analyzing soft such as BLAST [*J. Mol. Biol.*, 215, 403 (1990)], FASTA [*Methods in Enzymology*, 183, 63 (1990)] or the like.

Examples of the CHO cell of the present invention include a cell in which the enzyme activity is decreased or deleted.

rate, at a lectin concentration of at least 2 times, preferably 5 times, more preferably 10 times, and most preferably 20 times or more, than the parent cell line.

As the lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain, any lectin which can recognize the sugar chain structure can be used. Examples include a *Lens culinaris* lectin LCA (lentil agglutinin derived from *Lens culinaris*), a pea lectin PSA (pea lectin derived from *Pisum sativum*), a broad bean lectin VFA (agglutinin derived from *Vicia faba*), an *Aleuria aurantia* lectin AAL (lectin derived from *Aleuria aurantia*) and the like.

The CHO cell of the present invention can produce an antibody composition having higher ADCC activity than that of an antibody composition produced by the parent CHO cell before applying the technique for decreasing or deleting the enzyme activity of interest.

Also, the CHO cell of the present invention can produce an antibody composition having higher ADCC activity than that of an antibody composition in which, among the total complex *N*-glycoside-linked sugar chains bound to the Fc region contained in the antibody composition, the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain is less than 20%.

An example of the parent cell line to be used in the present invention is a cell in which the activity of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the activity of an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain is not decreased.

The present invention is described below in detail.

The gene disruption method may be any method, so long as it can disrupt the gene of the target enzyme is included. Examples include an antisense method, a ribozyme

method, a homologous recombination method, an RDO method, an RNAi method, a retrovirus-employed method, a transposon-employed method and the like. The methods are specifically described below.

(a) Preparation of the host cell of the present invention by the antisense method or the ribozyme method

The host cell of the present invention can be prepared by the ribozyme method described in *Cell Technology*, 12, 239 (1993); *BIO/TECHNOLOGY*, 17, 1097 (1999); *Hum. Mol. Genet.*, 5, 1083 (1995); *Cell Technology*, 13, 255 (1994); *Proc. Natl. Acad. Sci. USA*, 96, 1886 (1999); or the like, e.g., in the following manner by targeting at an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain.

A cDNA or a genome DNA encoding an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain is prepared.

The nucleotide sequence of the prepared cDNA or genome DNA is determined.

Based on the determined DNA sequence, an appropriate length of an antisense gene or ribozyme construct comprising a DNA moiety which encodes the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-

integrated into the chromosome and comprises a promoter at such a position that the designed antisense gene or ribozyme can be transferred is used. Examples include expression vectors which will be described later in the item 3.

Regarding the method for introducing a gene into various host cells, the methods for introducing recombinant vectors suitable for various host cells, which will be described later in the item 3, can be used.

The following method can be exemplified as the method for selecting a transformant as a measure of the activity of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the activity of an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain.

Method for selecting transformant:

Examples of the method for selecting a cell in which the activity of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the activity of an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain is decreased include biochemical methods or genetic engineering techniques described in *New Biochemical Experimentation Series 3-Saccharides I, Glycoprotein* (Tokyo Kagaku Dojin), edited by Japanese Biochemical society (1988); *Cell Engineering, Supplement, Experimental Protocol Series, Glycobiology Experimental Protocol, Glycoprotein, Glycolipid and Proteoglycan* (Shujun-sha), edited by Naoyuki Taniguchi, Akemi Suzuki, Kiyoshi Furukawa and Kazuyuki Sugawara (1996);, *Molecular Cloning, Second Edition*;

Current Protocols in Molecular Biology; and the like. Examples of the biochemical method include a method in which the enzyme activity is evaluated using an enzyme-specific substrate and the like. Examples of the genetic engineering technique include the Northern analysis, RT-PCR and the like which measures the amount of mRNA of a gene encoding the enzyme.

Examples of the method for selecting a transformant using the sugar chain structure of a glycoprotein on the cell membrane as a marker include the methods which will be described later in the item 1(5). Examples of the method for selecting a transformant using the sugar chain structure of a produced antibody molecule as a marker include the methods which will be described later in the items 5 and 6.

As the method for preparing cDNA encoding an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain, the following method is exemplified.

Preparation of DNA:

A total RNA or mRNA is prepared from a human or non-human animal tissue or cell.

A cDNA library is prepared from the prepared total RNA or mRNA.

Degenerative primers are produced based on the amino acid sequence of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain, and a gene fragment

encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of N-acetylglucosamine in the reducing end through α -bond in the complex N-glycoside-linked sugar chain is obtained by PCR using the prepared cDNA library as the template.

A DNA encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain can be obtained by screening the cDNA library using the obtained gene fragment as a probe.

Regarding the mRNA of a human or non-human tissue or cell, a commercially available product (e.g., manufactured by Clontech) may be used or it may be prepared from a human or non-human animal tissue or cell in the following manner. Examples of the method for preparing a total RNA from a human or non-human animal tissue or cell include the guanidine thiocyanate-caesium trifluoroacetate method [*Methods in Enzymology*, 154, 3 (1987)], the acidic guanidine thiocyanate phenol chloroform (AGPC) method [*Analytical Biochemistry*, 162, 156 (1987); *Experimental Medicine*, 9, 1937 (1991)] and the like.

Also, examples of the method for preparing mRNA from a total RNA as poly(A)⁺ RNA include an oligo(dT)-immobilized cellulose column method (*Molecular Cloning*, Second Edition) and the like.

In addition, mRNA can be prepared using a kit such as Fast Track mRNA Isolation Kit (manufactured by Invitrogen), Quick Prep mRNA Purification Kit (manufactured by Pharmacia) or the like.

A cDNA library is prepared from the prepared mRNA of a human or non-human animal tissue or cell. Examples of the method for preparing cDNA libraries include the methods described in *Molecular Cloning, Second Edition*; *Current Protocols in Molecular Biology; A Laboratory Manual, Second Edition* (1989); and the like, or methods using commercially available kits such as SuperScript Plasmid System for cDNA Synthesis and Plasmid Cloning (manufactured by Life Technologies), ZAP-cDNA Synthesis Kit (manufactured by STRATAGENE) and the like.

As the cloning vector for use in the preparation of the cDNA library, any vector such as a phage vector, a plasmid vector or the like can be used, so long as it is autonomously replicable in *Escherichia coli* K12. Examples include ZAP Express [manufactured by STRATAGENE, Strategies, 5, 58 (1992)], pBluescript II SK(+) [Nucleic Acids Research, 17, 9494 (1989)], Lambda ZAP II (manufactured by STRATAGENE), λ gt10 and λ gt11 [DNA Cloning, A Practical Approach, 1, 49 (1985)], λ TriplEx (manufactured by Clontech), λ ExCell (manufactured by Pharmacia), pcD2 [Mol. Cell. Biol., 3, 280 (1983)], pUC18 [Gene, 33, 103 (1985)] and the like.

Any microorganism can be used as the host microorganism, but *Escherichia coli* is preferably used. Examples include *Escherichia coli* XL1-Blue MRF' [manufactured by STRATAGENE, *Strategies*, 5, 81 (1992)], *Escherichia coli* C600 [Genetics, 39, 440 (1954)], *Escherichia coli* Y1088 [Science, 222, 778 (1983)], *Escherichia coli* Y1090 [Science, 222, 778 (1983)], *Escherichia coli* NM522 [J. Mol. Biol., 166, 1 (1983)], *Escherichia coli* K802 [J. Mol. Biol., 16, 118 (1966)], *Escherichia coli* JM105 [Gene, 38, 275 (1985)] and the like

The cDNA library may be used as such in the succeeding analysis, and in order to obtain a full length cDNA as efficient as possible by decreasing the ratio of an

infull length cDNA, a cDNA library prepared using the oligo cap method developed by Sugano et al. [Gene, 138, 171 (1994); Gene, 200, 149 (1997); Protein, Nucleic Acid and Protein, 41, 603 (1996); Experimental Medicine, 11, 2491 (1993); cDNA Cloning (Yodo-sha) (1996); Methods for Preparing Gene Libraries (Yodo-sha) (1994)] may be used in the following analysis.

Degenerative primers specific for the 5'-terminal and 3'-terminal nucleotide sequences of a nucleotide sequence presumed to encode the amino acid sequence are prepared based on the amino acid sequence of the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain, and DNA is amplified by PCR [PCR Protocols, Academic Press (1990)] using the prepared cDNA library as the template to obtain a gene fragment encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain.

It can be confirmed that the obtained gene fragment is a DNA encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain, by a method usually used for analyzing a nucleotide, such as the dideoxy method of Sanger et al. [Proc. Natl. Acad. Sci. USA, 74, 5463 (1977)], a nucleotide sequence analyzer such as ABIPRISM

377 DNA Sequencer (manufactured by PE Biosystems) or the like.

A DNA encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain can be obtained by carrying out colony hybridization or plaque hybridization (*Molecular Cloning*, Second Edition) for the cDNA or cDNA library synthesized from the mRNA contained in the human or non-human animal tissue or cell, using the gene fragment as a DNA probe.

Also, a DNA encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain can also be obtained by carrying out screening by PCR using the cDNA or cDNA library synthesized from the mRNA contained in a human or non-human animal tissue or cell as the template and using the primers used for obtaining the gene fragment encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain.

The nucleotide sequence of the obtained DNA encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in

the complex *N*-glycoside-linked sugar chain is analyzed from its terminus and determined by a method usually used for analyzing a nucleotide, such as the dideoxy method of Sanger et al. [*Proc. Natl. Acad. Sci. USA*, 74, 5463 (1977)], a nucleotide sequence analyzer such as ABIPRISM 377 DNA Sequencer (manufactured by PE Biosystems) or the like.

A gene encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain can also be determined from genes in data bases by searching nucleotide sequence data bases such as GenBank, EMBL, DDBJ and the like using a homology retrieving program such as BLAST based on the determined cDNA nucleotide sequence.

Examples of the nucleotide sequence of the gene obtained by the method encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose include the nucleotide sequence represented by SEQ ID NO:48, 51 or 65. Examples of the nucleotide sequence of the gene encoding the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain include the nucleotide sequence represented by SEQ ID NO:1 or 2.

The cDNA encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain can also be obtained by chemically synthesizing it with a DNA synthesizer such as DNA Synthesizer model 392 manufactured by Perkin Elmer

- 61 -

chain include the nucleotide sequence represented by SEQ ID NO:3.

In addition, the host cell of the present invention can also be obtained without using an expression vector, by directly introducing an antisense oligonucleotide or ribozyme into a host cell, which is designed based on the nucleotide sequence encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain.

The antisense oligonucleotide or ribozyme can be prepared in the usual method or using a DNA synthesizer. Specifically, it can be prepared based on the sequence information of an oligonucleotide having a corresponding sequence of continued 5 to 150 bases, preferably 5 to 60 bases, and more preferably 10 to 40 bases, among nucleotide sequences of a cDNA and a genome DNA encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain, by synthesizing an oligonucleotide which corresponds to a sequence complementary to the oligonucleotide (antisense oligonucleotide) or a ribozyme comprising the oligonucleotide sequence.

Examples of the oligonucleotide include oligo RNA and derivatives of the oligonucleotide (hereinafter referred to as "oligonucleotide derivatives").

Examples of the oligonucleotide derivatives include oligonucleotide derivatives in which a phosphodiester bond in the oligonucleotide is converted into a phosphorothioate bond, an oligonucleotide derivative in which a

- 63 -

University Press (1993); *Biomanual Series 8, Gene Targeting, Preparation of Mutant Mice using ES Cells*, Yodo-sha (1995) (hereinafter referred to as "*Preparation of Mutant Mice using ES Cells*"); or the like, for example, as follows.

A genome DNA encoding an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain is prepared.

Based on the nucleotide sequence of the genome DNA, a target vector is prepared for homologous recombination of a target gene to be modified (e.g., structural gene of the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain, or a promoter gene).

The host cell of the present invention can be produced by introducing the prepared target vector into a host cell and selecting a cell in which homologous recombination occurred between the target gene and target vector.

As the host cell, any cell such as yeast, animal cell, insect cell or plant cell can be used, so long as it has a gene encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain. Examples include the host cells which will be described later in the item 3.

Approach, IRL Press at Oxford University Press (1993); *Biomanual Series 8, Gene Targeting, Preparation of Mutant Mice using ES Cells*, Yodo-sha (1995); or the like. Examples of the method for selecting the homologous recombinant of interest from the selected cell lines include the Southern hybridization method for genome DNA (*Molecular Cloning*, Second Edition), PCR [*PCR Protocols*, Academic Press (1990)], and the like.

(c) Preparation of the host cell of the present invention by RDO method

The host cell of the present invention can be prepared by an RDO (RNA-DNA oligonucleotide) method by targeting at a gene encoding an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain, for example, as follows

A cDNA or a genome DNA encoding an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain is prepared.

The nucleotide sequence of the prepared cDNA or genome DNA is determined.

Based on the determined DNA sequence, an appropriate length of an RDO construct comprising a DNA moiety which encodes the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in

the complex *N*-glycoside-linked sugar chain or a part of its non-translation region or an intron, is designed and synthesized.

The host cell of the present invention can be obtained by introducing the synthesized RDO into a host cell and then selecting a transformant in which a mutation occurred in the target enzyme, namely the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain.

As the host cell, any cell such as yeast, animal cell, insect cell or plant cell can be used, so long as it has a gene encoding the target enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or of the target enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain. Examples include the host cells which will be described later in the item 3.

Examples of the method for introducing RDO into various host cells include the methods for introducing recombinant vectors suited for various host cells, which will be described later in the item 3.

Examples of the method for preparing cDNA encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain include the methods described in the preparation of DNA in the item 1(1)(a) and the like.

Examples of the method for preparing a genome DNA encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain include the methods in preparation of genome DNA described in the item 1(1)(a) and the like.

The nucleotide sequence of the DNA can be determined by digesting it with appropriate restriction enzymes, cloning the fragments into a plasmid such as pBluescript SK(-) (manufactured by Stratagene) or the like, subjecting the clones to the reaction generally used as a method for analyzing a nucleotide sequence such as the dideoxy method [Proc. Natl. Acad. Sci. USA, 74, 5463 (1977)] of Sanger et al. or the like, and then analyzing the clones using an automatic nucleotide sequence analyzer such as A.L.F. DNA Sequencer (manufactured by Pharmacia) or the like.

The RDO can be prepared by a usual method or using a DNA synthesizer.

Examples of the method for selecting a cell in which a mutation occurred, by introducing the ROD into the host cell, in the gene encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain include the methods for directly detecting mutations in chromosomal genes described in *Molecular Cloning*, Second Edition, *Current Protocols in Molecular Biology* and the like;

the methods described in the item 1(1)(a) for selecting a transformant through the evaluation of the

activity of the introduced enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of N-acetylglucosamine in the reducing end through α -bond in the complex N-glycoside-linked sugar chain; the method for selecting a transformant using the sugar structure of a glycoprotein on the cell membrane which will be described later in the item 1(5); and the method for selecting a transformant as a measure of the sugar structure of the produced antibody molecule which will be described later in the item 5 or 6, and the like.

The construct of the ROD can be designed in accordance with the methods described in *Science*, 273, 1386 (1996); *Nature Medicine*, 4, 285 (1998); *Hepatology*, 25, 1462 (1997); *Gene Therapy*, 5, 1960 (1999); *J. Mol. Med.*, 75, 829 (1997); *Proc. Natl. Acad. Sci. USA*, 96, 8774 (1999); *Proc. Natl. Acad. Sci. USA*, 96, 8768 (1999); *Nuc. Acids. Res.*, 27, 1323 (1999); *Invest. Dermatol.*, 111, 1172 (1998); *Nature Biotech.*, 16, 1343 (1998); *Nature Biotech.*, 18, 43 (2000); *Nature Biotech.*, 18, 555 (2000); and the like.

(d) Preparation of the host cell of the present invention by RNAi method

The host cell of the present invention can be prepared by the RNAi (RNA interference) method by targeting at a gene of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or of an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of N-acetylglucosamine in the reducing end through α -bond in the complex N-glycoside-linked sugar chain, for example, as follows.

A cDNA encoding an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or an

enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain is prepared.

The nucleotide sequence of the prepared cDNA is determined.

Based on the determined DNA sequence, an appropriate length of an RNAi gene construct comprising the DNA coding moiety encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain or a part of its non-translation region, is designed.

In order to express the RNAi gene in a cell, a recombinant vector is prepared by inserting a fragment or full length of the prepared DNA into downstream of the promoter of an appropriate expression vector.

A transformant is obtained by introducing the recombinant vector into a host cell suitable for the expression vector.

The host cell of the present invention can be obtained by selecting a transformant as a measure of the activity of the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain, or the sugar chain structure of a glycoprotein on the cell membrane or of the produced antibody molecule.

As the host cell, any cell such as yeast, animal cell, insect cell or plant cell can be used, so long as it has a gene encoding the target enzyme relating to the

synthesis of an intracellular sugar nucleotide, GDP-fucose or the target enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain. Examples include the host cells which will be described later in the item 3.

As the expression vector, a vector which is autonomously replicable in the host cell or can be integrated into the chromosome and comprises a promoter at such a position that the designed RNAi gene can be transferred is used. Examples include the expression vectors which will be described later in the item 3.

As the method for introducing a gene into various host cells, the methods for introducing recombinant vectors suitable for various host cells, which will be described later in the item 3, can be used.

Examples of the method for selecting a transformant as a measure of the activity of the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain include the methods described in the item 1(1)(a).

Examples of the method for selecting a transformant as a measure of the sugar chain structure of a glycoprotein on the cell membrane include the methods which will be described later in the item 1(5). Examples of the method for selecting a transformant as a measure of the sugar chain structure of a produced antibody molecule include the methods which will be described later in the item 5 or 6.

Examples of the method for preparing cDNA encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the

linked sugar chain, or the sugar chain structure of a glycoprotein of a produced antibody molecule or on the cell membrane.

The transposon system is a system in which a mutation is induced by randomly inserting an exogenous gene into chromosome, wherein an exogenous gene interposed between transposons is generally used as a vector for inducing a mutation, and a transposase expression vector for randomly inserting the gene into chromosome is introduced into the cell at the same time.

Any transposase can be used, so long as it is suitable for the sequence of the transposon to be used.

As the exogenous gene, any gene can be used, so long as it can induce a mutation in the DNA of a host cell.

As the host cell, any cell such as yeast, animal cell, insect cell or plant cell can be used, so long as it has a gene encoding the target enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or of the target enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain. Examples include the host cells which will be described later in the item 3. For introducing the gene into various host cells, the method for introducing recombinant vectors suitable for various host cells, which will be described later in the item 3, can be used.

Examples of the method for selecting a mutant as a measure of the activity of the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain include the methods described in the item 1(1)(a).

- 74 -

- 77 -

Examples of the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose include GMD, Fx, GFPP, fucokinase and the like. Examples of the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain include α -1,6-fucosyltransferase, α -L-fucosidase and the like.

Examples of the method include 1) a method in which a desired cell line is selected from mutants obtained by a mutation-inducing treatment of a parent cell line with a mutagen or spontaneously generated mutants, as a measure of the activity of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the activity of an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain, 2) a method in which a desired cell line is selected from mutants obtained by a mutation-inducing treatment of a parent cell line with a mutagen or spontaneously generated mutants, as a measure of the sugar chain structure of a produced antibody molecule and 3) a method in which a desired cell line is selected from mutants obtained by a mutation-inducing treatment of a parent cell line with a mutagen or spontaneously generated mutants, as a measure of the sugar chain structure of a glycoprotein on the cell membrane.

As the mutation-inducing treatment, any treatment can be used, so long as it can induce a point mutation or a deletion or frame shift mutation in the DNA of cells of the parent cell line.

Examples include treatment with ethyl nitrosourea, nitrosoguanidine, benzopyrene or an acridine pigment and treatment with radiation. Also, various alkylating agents and carcinogens can be used as mutagens. Examples of the

method for allowing a mutagen to act upon cells include the methods described in *Tissue Culture Techniques*, 3rd edition (Asakura Shoten), edited by Japanese Tissue Culture Association (1996), *Nature Genet.*, 24, 314 (2000) and the like.

Examples of the spontaneously generated mutant include mutants which are spontaneously formed by continuing subculture under general cell culture conditions without applying special mutation-inducing treatment.

Examples of the method for measuring the activity of the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the activity of the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain include the methods described in the item 1(1)(a). Examples of the method for discriminating the sugar chain structure of a prepared antibody molecule include the methods which will be described later in the item 5 or 6. Examples of the method for discriminating the sugar chain structure of a glycoprotein on the cell membrane include the methods which will be described later in the item 1(5).

(4) Method for inhibiting transcription and/or translation of a gene encoding an enzyme

The host cell of the present invention can be prepared by inhibiting transcription and/or translation of a target gene through a method such as the antisense RNA/DNA technique [*Bioscience and Industry*, 50, 322 (1992); *Chemistry*, 46, 681 (1991); *Biotechnology*, 2, 358 (1992); *Trends in Biotechnology*, 10, 87 (1992); *Trends in Biotechnology*, 10, 152 (1992); *Cell Engineering*, 16, 1463 (1997)], the triple helix technique [*Trends in Biotechnology*, 10, 132 (1992)] or the like, using a gene

encoding an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose and/or an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain, as the target.

Examples of the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose include GMD, Fx, GFPP, fucokinase and the like. Examples of the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain include α -1,6-fucosyltransferase, α -L-fucosidase and the like.

(5) Method for selecting a cell line resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain

The host cell of the present invention can be prepared by using a method for selecting a cell line resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain.

Examples of the method for selecting a cell line resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain include the methods using lectin described in *Somatic Cell Mol. Genet.*, 12, 51 (1986) and the like. As the lectin, any lectin can be used, so long as it is a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to

- 81 -

chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain, as the target.

In a transgenic non-human animal, the embryonic stem cell of the present invention in which the activity of the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the activity of the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain is controlled can be prepared applying the method similar to that in the item 1 to an embryonic stem cell of the intended non-human animal such as cattle, sheep, goat, pig, horse, mouse, rat, fowl, monkey, rabbit or the like.

Specifically, a mutant clone is prepared in which a gene encoding the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain is inactivated or substituted with any sequence, by a known homologous recombination technique [e.g., *Nature*, 326, 6110, 295 (1987); *Cell*, 51, 3, 503 (1987); or the like]. Using the prepared mutant clone, a chimeric individual comprising an embryonic stem cell clone and a normal cell can be prepared by an injection chimera method into blastocyst of fertilized egg of an animal or by an aggregation chimera method. The chimeric individual is crossed with a normal individual, so that a transgenic non-human animal in which the activity of the enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the activity of the enzyme relating to the modification of a sugar chain

wherein 1-position of fucose is bound to 6-position of N-acetylglucosamine in the reducing end through α -bond in the complex N-glycoside-linked sugar chain is decreased or deleted in the whole body cells can be obtained.

Also, a fertilized egg cell of the present invention in which the activity of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the activity of an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of N-acetylglucosamine in the reducing end through α -bond in the complex N-glycoside-linked sugar chain is decreased or deleted can be prepared by applying the method similar to that in the item 1 to fertilized egg of a non-human animal of interest such as cattle, sheep, goat, pig, horse, mouse, rat, fowl, monkey, rabbit or the like.

A transgenic non-human animal in which the activity of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the activity of an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of N-acetylglucosamine in the reducing end through α -bond in the complex N-glycoside-linked sugar chain is decreased can be prepared by transplanting the prepared fertilized egg cell into the oviduct or uterus of a pseudopregnant female using the embryo transplantation method described in *Manipulating Mouse Embryo*, Second Edition or the like, followed by childbirth by the animal.

In a transgenic plant, the callus of the present invention in which the activity of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the activity of an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of N-acetylglucosamine in the reducing end through α -bond in the complex N-glycoside-

A transgenic plant in which the activity of an enzyme relating to the synthesis of an intracellular sugar nucleotide, GDP-fucose or the activity of an enzyme relating to the modification of a sugar chain wherein 1-position of fucose is bound to 6-position of N-acetylglucosamine in the reducing end through α -bond in the complex N-glycoside-linked sugar chain is decreased can be prepared by culturing the prepared callus using a medium comprising auxin and cytokinin to redifferentiate it in accordance with a known method [Tissue Culture, 20 (1994); Tissue Culture, 21 (1995); Trends in Biotechnology, 15, 45 (1997)].

3. Method for producing an antibody composition

The antibody composition can be obtained by expressing it in a host cell using the methods described in *Molecular Cloning*, Second Edition; *Current Protocols in Molecular Biology*; *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988 (hereinafter referred also to as "Antibodies"); *Monoclonal Antibodies: Principles and Practice*, Third Edition, Acad. Press, 1993 (hereinafter referred also to as "Monoclonal Antibodies"); and *Antibody Engineering, A Practical Approach*, IRL Press at Oxford University Press (hereinafter referred also to as "Antibody Engineering"), for example, as follows.

A full length cDNA encoding an antibody molecule is prepared, and an appropriate length of a DNA fragment comprising a moiety encoding the antibody molecule is prepared.

A recombinant vector is prepared by inserting the DNA fragment or the full length cDNA into downstream of the promoter of an appropriate expression vector.

A transformant which produces the antibody molecule can be obtained by introducing the recombinant vector into a host cell suitable for the expression vector.

As the host cell, any of yeast, animal cell, insect cell, plant cell or the like can be used, so long as it can express the gene of interest.

A cell such as yeast, animal cell, insect cell, plant cell or the like into which an enzyme relating to the modification of an *N*-glycoside-linked sugar chain which binds to the Fc region of the antibody molecule is introduced by a genetic engineering technique can also be used as the host cell.

As the expression vector, a vector which is autonomously replicable in the host cell or can be integrated into the chromosome and comprises a promoter at such a position that the DNA encoding the antibody molecule of interest can be transferred is used.

The cDNA can be prepared from a human or non-human tissue or cell using, e.g., a probe primer specific for the antibody molecule of interest, in accordance with the methods described in the preparation of DNA in the item 1(1)(a).

When a yeast is used as the host cell, examples of the expression vector include YEP13 (ATCC 37115), YEp24 (ATCC 37051), YCp50 (ATCC 37419) and the like.

Any promoter can be used, so long as it can function in yeast. Examples include a promoter of a gene of the glycolytic pathway such as a hexose kinase gene, etc., PH05 promoter, PGK promoter, GAP promoter, ADH promoter, gal 1 promoter, gal 10 promoter, heat shock protein promoter, MF α 1 promoter, CUP 1 promoter and the like.

Examples of the host cell include microorganisms belonging to the genus *Saccharomyces*, the genus *Schizosaccharomyces*, the genus *Kluyveromyces*, the genus

Trichosporon, the genus *Schwanniomyces* and the like, such as *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Trichosporon pullulans* and *Schwanniomyces alluvius*, etc.

As the method for introducing the recombinant vector, any method can be used, so long as it can introduce DNA into yeast. Examples include electroporation [*Methods in Enzymology*, 194, 182 (1990)], spheroplast method [*Proc. Natl. Acad. Sci. USA*, 84, 1929 (1978)], lithium acetate method [*J. Bacteriol.*, 153, 163 (1983)], a method described in *Proc. Natl. Acad. Sci. USA*, 75, 1929 (1978) and the like.

When an animal cell is used as the host, examples of the expression vector include pcDNA1, pCDM8 (available from Funakoshi), pAGE107 [Japanese Published Examined Patent Application No. 22979/91; *Cytotechnology*, 3, 133 (1990)], pAS3-3 (Japanese Published Examined Patent Application No. 227075/90), pCDM8 [*Nature*, 329, 840 (1987)], pcDNA1/Amp (manufactured by Invitrogen), pREP4 (manufactured by Invitrogen), pAGE103 [*J. Biochemistry*, 101, 1307 (1987)], pAGE210 and the like.

Any promoter can be used, so long as it can function in an animal cell. Examples include a promoter of IE (immediate early) gene of cytomegalovirus (CMV), an early promoter of SV40, a promoter of retrovirus, a promoter of metallothionein, a heat shock promoter, an SR α promoter and the like. Also, an enhancer of the IE gene of human CMV may be used together with the promoter.

Examples of the host cell include a human cell such as Namalwa cell, a monkey cell such as COS cell, a Chinese hamster cell such as CHO cell or HBT5637 (Japanese Published Examined Patent Application No. 299/88), a rat myeloma cell, a mouse myeloma cell, a cell derived from syrian hamster kidney, an embryonic stem cell, a fertilized egg cell and the like.

(1992)], a *Trichoplusia ni* oocyte High 5 (manufactured by Invitrogen) and the like.

Examples of the method for the simultaneously introducing the recombinant gene-introducing vector and the baculovirus for preparing the recombinant virus include the calcium phosphate method (Japanese Published Examined Patent Application No. 227075/90), the lipofection method [Proc. Natl. Acad. Sci. USA, 84, 7413 (1987)] and the like.

When a plant cell is used as the host, examples of the expression vector include Ti plasmid, tobacco mosaic virus and the like.

As the promoter, any promoter can be used, so long as it can function in a plant cell. Examples include cauliflower mosaic virus (CaMV) 35S promoter, rice actin 1 promoter and the like.

Examples of the host cell include plant cells of tobacco, potato, tomato, carrot, soybean, rape, alfalfa, rice, wheat, barley, etc., and the like.

As the method for introducing the recombinant vector, any method can be used, so long as it can introduce DNA into a plant cell. Examples include a method using *Agrobacterium* (Japanese Published Examined Patent Application No. 140885/84, Japanese Published Examined Patent Application No. 70080/85, WO 94/00977), electroporation (Japanese Published Examined Patent Application No. 251887/85), a method using a particle gun (gene gun) (Japanese Patent No. 2606856, Japanese Patent No. 2517813) and the like.

As the method for expressing a gene, secretion production, expression of a fusion protein of the Fc region with other protein and the like can be carried out in accordance with the method described in *Molecular Cloning*, Second Edition or the like, in addition to the direct expression.

When a gene is expressed by a bacterium, a yeast, an animal cell, an insect cell or a plant cell into which a gene relating to the synthesis of a sugar chain is introduced, an antibody molecule to which a sugar or a sugar chain is added by the introduced gene can be obtained.

An antibody composition can be obtained by culturing the obtained transformant in a medium to produce and accumulate the antibody molecule in the culture and then recovering it from the resulting culture. The method for culturing the transformant using a medium can be carried out in accordance with a general method which is used for the culturing of host cells.

As the medium for culturing a transformant obtained using a prokaryote such as *Escherichia coli* etc. or a eukaryote such as yeast etc. as the host cell, the medium may be either a natural medium or a synthetic medium, so long as it comprises materials such as a carbon source, a nitrogen source, an inorganic salt and the like which can be assimilated by the organism and culturing of the transformant can be efficiently carried out.

As the carbon source, those which can be assimilated by the organism can be used. Examples include carbohydrates such as glucose, fructose, sucrose, molasses containing them, starch, starch hydrolysate, etc.; organic acids such as acetic acid, propionic acid, etc.; alcohols such as ethanol, propanol, etc.; and the like.

Examples of the nitrogen source include ammonia; ammonium salts of inorganic acid or organic acid such as ammonium chloride, ammonium sulfate, ammonium acetate, ammonium phosphate, etc.; other nitrogen-containing compounds; peptone; meat extract; yeast extract; corn steep liquor; casein hydrolysate; soybean meal; soybean meal hydrolysate; various fermented cells and hydrolysates thereof; and the like.

Examples of the inorganic material include potassium dihydrogen phosphate, dipotassium hydrogen phosphate, magnesium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, copper sulfate, calcium carbonate, and the like

The culturing is carried out generally under aerobic conditions such as a shaking culture, submerged-aeration stirring culture or the like. The culturing temperature is preferably 15 to 40°C, and the culturing time is generally 16 hours to 7 days. During the culturing, the pH is maintained at 3.0 to 9.0. The pH is adjusted using an inorganic or organic acid, an alkali solution, urea, calcium carbonate, ammonia or the like.

If necessary, an antibiotic such as ampicillin, tetracycline or the like may be added to the medium during the culturing.

When a microorganism transformed with a recombinant vector obtained using an inducible promoter as the promoter is cultured, an inducer may be added to the medium, if necessary. For example, when a microorganism transformed with a recombinant vector obtained using lac promoter is cultured, isopropyl- β -D-thiogalactopyranoside may be added to the medium, and when a microorganism transformed with a recombinant vector obtained using trp promoter is cultured, indoleacrylic acid may be added to the medium.

When a transformant obtained using an animal cell as the host cell is cultured, examples of the medium include generally used RPMI 1640 medium [*The Journal of the American Medical Association*, 199, 519 (1967)], Eagle's MEM medium [*Science*, 122, 501 (1952)], Dulbecco's modified MEM medium [*Virology*, 8, 396 (1959)], 199 medium [*Proceeding of the Society for the Biological Medicine*, 73, 1 (1950)] and Whitten's medium [*Developmental Engineering Experimentation Manual-Preparation of Transgenic Mice* (Kodan-sha), edited

antibody molecule is inserted, in accordance with a general culturing method, to thereby produce and accumulate the antibody composition, and then recovering the antibody composition from the culture.

As the method for expressing the gene, secretion production, expression of a fusion protein and the like can be carried out in accordance with the method described in *Molecular Cloning*, Second Edition, in addition to the direct expression.

Examples of the method for producing an antibody composition include a method of intracellular expression in a host cell, a method of extracellular secretion from a host cell, and a method of production on a host cell membrane outer envelope. The method can be selected by changing the host cell used or the structure of the antibody composition produced.

When the antibody composition of the present invention is produced in a host cell or on a host cell membrane outer envelope, it can be positively secreted extracellularly in accordance with the method of Paulson et al. [*J. Biol. Chem.*, 264, 17619 (1989)], the method of Lowe et al. [*Proc. Natl. Acad. Sci. USA*, 86, 8227 (1989), *Genes Develop.*, 4, 1288 (1990)], the methods described in Japanese Published Examined Patent Application No. 336963/93 and Japanese Published Examined Patent Application No. 823021/94 and the like.

That is, an antibody molecule of interest can be positively secreted extracellularly from a host cell by inserting a DNA encoding the antibody molecule and a DNA encoding a signal peptide suitable for the expression of the antibody molecule into an expression vector using a gene recombination technique, introducing the expression vector into the host cell and then expressing the antibody molecule.

In the case of an animal individual, an antibody composition can be produced by rearing a transgenic non-human animal into which a DNA encoding an antibody molecule is introduced to thereby produce and accumulate the antibody composition in the animal, and then recovering the antibody composition from the animal. Examples of the place of the animal where the composition is produced and accumulated include milk (Japanese Published Examined Patent Application No. 309192/88) and eggs of the animal. As the promoter used in this case, any promoter can be used, so long as it can function in an animal. Preferred examples include mammary gland cell-specific promoters such

as α casein promoter, β casein promoter, β lactoglobulin promoter, whey acidic protein promoter and the like.

Example of the method for producing an antibody composition using a plant individual include a method in which an antibody composition is produced by cultivating a transgenic plant into which a DNA encoding an antibody molecule is introduced by a known method [*Tissue Culture*, 20 (1994); *Tissue Culture*, 21 (1995); *Trends in Biotechnology*, 15, 45 (1997)] to produce and accumulate the antibody composition in the plant, and then recovering the antibody composition from the plant.

Regarding purification of an antibody composition produced by a transformant into which a gene encoding an antibody molecule is introduced, for example, when the antibody composition is intracellularly expressed in a dissolved state, the cells after culturing are recovered by centrifugation, suspended in an aqueous buffer and then disrupted using ultrasonic oscillator, French press, Manton Gaulin homogenizer, dynamill or the like to obtain a cell-free extract. A purified product of the antibody composition can be obtained from a supernatant obtained by centrifuging the cell-free extract, by using a general enzyme isolation purification techniques such as solvent extraction; salting out; desalting with ammonium sulfate, etc.; precipitation with an organic solvent; anion exchange chromatography using a resin such as diethylaminoethyl (DEAE)-Sephadex, DIAION HPA-75 (manufactured by Mitsubishi Chemical), etc.; cation exchange chromatography using a resin such as S-Sephadex FF (manufactured by Pharmacia), etc.; hydrophobic chromatography using a resin such as butyl-Sephadex, phenyl-Sephadex, etc.; gel filtration using a molecular sieve; affinity chromatography; chromatofocusing; electrophoresis such as isoelectric focusing, etc.; and the like which may be used alone or in combination.

The humanized antibody expression vector may be either of a type in which genes encoding the H chain and L chain of an antibody exist on separate vectors or of a type in which both genes exist on the same vector (tandem type). In respect of easiness of construction of a humanized antibody expression vector, easiness of introduction into animal cells, and balance between the expression amounts of the H and L chains of an antibody in animal cells, a tandem

The constructed humanized antibody expression vector can be used for expression of a human chimeric antibody and a human CDR-grafted antibody in animal cells.

cdNAs encoding the H chain and L chain V regions of an antibody derived from an animal other than human, such as a mouse antibody, can be obtained in the following manner.

As the animal other than human, any animal such as mouse, rat, hamster, rabbit or the like can be used so long as a hybridoma cell can be produced therefrom.

- 97 -

cellulose column method (*Molecular Cloning*, Second Edition) and the like. In addition, examples of a kit for preparing mRNA from a hybridoma cell include Fast Track mRNA Isolation Kit (manufactured by Invitrogen), Quick Prep mRNA Purification Kit (manufactured by Pharmacia) and the like.

Examples of the method for synthesizing cDNA and preparing a cDNA library include the usual methods (*Molecular Cloning*, Second Edition, *Current Protocols in Molecular Biology*, Supplement 1-34), methods using a commercially available kit such as SuperScript™, Plasmid System for cDNA Synthesis and Plasmid Cloning (manufactured by GIBCO BRL) or ZAP-cDNA Synthesis Kit (manufactured by Stratagene), and the like.

In preparing the cDNA library, the vector into which a cDNA synthesized using mRNA extracted from a hybridoma cell as the template is inserted may be any vector so long as the cDNA can be inserted. Examples include ZAP Express [*Strategies*, 5, 58 (1992)], pBluescript II SK(+) [*Nucleic Acids Research*, 17, 9494 (1989)], λzapII (manufactured by Stratagene), λgt10 and λgt11 [*DNA Cloning, A Practical Approach*, I, 49 (1985)], Lambda BlueMid (manufactured by Clontech), λExCell, pT7T3 18U (manufactured by Pharmacia), pCD2 [*Mol. Cell. Biol.*, 3, 280 (1983)], pUC18 [*Gene*, 33, 103 (1985)] and the like.

As *Escherichia coli* into which the cDNA library constructed from a phage or plasmid vector is introduced, any *Escherichia coli* can be used, so long as the cDNA library can be introduced, expressed and maintained. Examples include XL1-Blue MRF' [*Strategies*, 5, 81 (1992)], C600 [*Genetics*, 39, 440 (1954)], Y1088 and Y1090 [*Science*, 222, 778 (1983)], NM522 [*J. Mol. Biol.*, 166, 1 (1983)], K802 [*J. Mol. Biol.*, 16, 118 (1966)], JM105 [*Gene*, 38, 275 (1985)] and the like.

As the method for selecting a cDNA clone encoding the H chain and L chain V regions of an antibody derived

from an animal other than human from the cDNA library, a colony hybridization or a plaque hybridization using an isotope- or fluorescence-labeled probe can be used (*Molecular Cloning*, Second Edition). The cDNA encoding the H chain and L chain V regions can also be prepared by preparing primers and carrying out polymerase chain reaction (hereinafter referred to as "PCR"; *Molecular Cloning*, Second Edition; *Current Protocols in Molecular Biology*, Supplement 1-34) using a cDNA synthesized from mRNA or a cDNA library as the template.

The nucleotide sequences of the cDNAs can be determined by digesting the selected cDNAs with appropriate restriction enzymes, cloning the fragments into a plasmid such as pBluescript SK(-) (manufactured by Stratagene) or the like, carrying out the reaction of a generally used nucleotide sequence analyzing method such as the dideoxy method [*Proc. Natl. Acad. Sci. USA*, 74, 5463 (1977)] of Sanger et al. or the like and then analyzing the clones using an automatic nucleotide sequence analyzer such as A.L.F. DNA Sequencer (manufactured by Pharmacia) or the like. Whether or not the obtained cDNAs are encoding the full amino acid sequences of the H chain and L chain V regions of the antibody containing a secretory signal sequence can be confirmed by deducing the full amino acid sequences of the H chain and L chain V regions from the determined nucleotide sequence and comparing them with the full amino acid sequences of the H chain and L chain V regions of known antibodies [*Sequences of Proteins of Immunological Interest*, US Dep. Health and Human Services (1991)].

(3) Analysis of amino acid sequence of V region of antibody derived from animal other than human

Regarding the full amino acid sequences of the H chain and L chain V regions of the antibody containing a

secretory signal sequence, the length of the secretory signal sequence and the N-terminal amino acid sequences can be deduced and subgroups to which they belong can also be found, by comparing them with the full amino acid sequences of the H chain and L chain V regions of known antibodies [*Sequences of Proteins of Immunological Interest*, US Dep. Health and Human Services, (1991)]. In addition, the amino acid sequences of the H chain and L chain V regions of each CDR can also be found by comparing them with the amino acid sequences of the H chain and L chain V regions of known antibodies [*Sequences of Proteins of Immunological Interest*, US Dep. Health and Human Services, (1991)].

(4) Construction of human chimeric antibody expression vector

A human chimeric antibody expression vector can be constructed by cloning cDNAs encoding the H chain and L chain V regions of an antibody derived from an animal other than human into upstream of genes encoding the H chain and L chain C regions of a human antibody in the vector for humanized antibody expression constructed in the item 3(1). For example, a human chimeric antibody expression vector can be constructed by linking each of cDNAs encoding the H chain and L chain V regions of an antibody derived from an animal other than human to a synthetic DNA comprising nucleotide sequences at the 3'-terminals of the H chain and L chain V regions of an antibody derived from an animal other than human and nucleotide sequences at the 5'-terminals of the H chain and L chain C regions of a human antibody and also having a recognition sequence of an appropriate restriction enzyme at both terminals, and by cloning them into upstream of genes encoding the H chain and L chain C regions of a human antibody contained in the vector for humanized antibody expression constructed described in the item 3(1).

(5) Construction of cDNA encoding V region of human CDR-grafted antibody

cDNAs encoding the H chain and L chain V regions of a human CDR-grafted antibody can be obtained as follows. First, amino acid sequences of the frameworks (hereinafter referred to as "FR") of the H chain and L chain V regions of a human antibody for grafting CDR of the H chain and L chain V regions of an antibody derived from an animal other than human is selected. As the amino acid sequences of FRs of the H chain and L chain V regions of a human antibody, any amino acid sequences can be used so long as they are derived from a human antibody. Examples include amino acid sequences of FRs of the H chain and L chain V regions of human antibodies registered at databases such as Protein Data Bank, etc., amino acid sequences common in each subgroup of FRs of the H chain and L chain V regions of human antibodies [*Sequences of Proteins of Immunological Interest*, US Dep. Health and Human Services (1991)] and the like. But in order to produce a human CDR-grafted antibody having potent activity, it is preferable to select an amino acid sequence having a homology as high as possible (at least 60% or more) with amino acid sequences of the H chain and L chain V regions of an antibody of interest derived from an animal other than human.

Next, the amino acid sequences of CDRs of the H chain and L chain V regions of the antibody of interest derived from an animal other than human are grafted to the selected amino acid sequences of FRs of the H chain and L chain V regions of a human antibody to design amino acid sequences of the H chain and L chain V regions of the human CDR-grafted antibody. The designed amino acid sequences are converted into DNA sequences by considering the frequency of codon usage found in nucleotide sequences of antibody genes [*Sequences of Proteins of Immunological*

Interest, US Dep. Health and Human Services (1991)], and the DNA sequences encoding the amino acid sequences of the H chain and L chain V regions of the human CDR-grafted antibody are designed. Based on the designed DNA sequences, several synthetic DNA fragments having a length of about 100 bases are synthesized, and PCR is carried out using them. In this case, it is preferable in each of the H chain and the L chain that 6 synthetic DNAs are designed in view of the reaction efficiency of PCR and the lengths of DNAs which can be synthesized.

Also, they can be easily cloned into the vector for humanized antibody expression constructed in the item 3(1) by introducing recognition sequences of an appropriate restriction enzyme into the 5'-terminals of the synthetic DNA present on both terminals. After the PCR, the amplified product is cloned into a plasmid such as pBluescript SK(-) (manufactured by Stratagene) or the like and the nucleotide sequences are determined by the method in the item 3(2) to thereby obtain a plasmid having DNA sequences encoding the amino acid sequences of the H chain and L chain V regions of the desired human CDR-grafted antibody.

(6) Construction of human CDR-grafted antibody expression vector

A human CDR-grafted antibody expression vector can be constructed by cloning the cDNAs encoding the H chain and L chain V regions of the human CDR-grafted antibody constructed in the item 3(5) into upstream of the gene encoding H chain and L chain C regions of a human antibody in the vector for humanized antibody expression described in the item 3(1). For example, the human CDR-grafted antibody expression vector can be constructed by introducing recognizing sequences of an appropriate restriction enzyme into the 5'-terminals of both terminals

of a synthetic DNA fragment, among the synthetic DNA fragments which are used when PCR is carried out in the item 3(5) for constructing the H chain and L chain V regions of the human CDR-grafted antibody, so that they are cloned into upstream of the genes encoding the H chain and L chain C regions of a human antibody in the vector for humanized antibody expression described in the item 3(1) in such a manner that they can be expressed in a suitable form.

(7) Stable production of humanized antibody

A transformant capable of stably producing a human chimeric antibody and a human CDR-grafted antibody (both hereinafter referred to as "humanized antibody") can be obtained by introducing the humanized antibody expression vectors described in the items 3(4) and (6) into an appropriate animal cell.

Examples of the method for introducing a humanized antibody expression vector into an animal cell include electroporation [Japanese Published Examined Patent Application No. 257891/90, *Cytotechnology*, 3, 133 (1990)] and the like.

As the animal cell into which a humanized antibody expression vector is introduced, any cell can be used so long as it is an animal cell which can produce the humanized antibody.

Examples include mouse myeloma cells such as NSO cell and SP2/0 cell, Chinese hamster ovary cells such as CHO/dhfr⁻ cell and CHO/DG44 cell, rat myeloma such as YB2/0 cell and IR983F cell, BHK cell derived from a syrian hamster kidney, a human myeloma cell such as Namalwa cell, and the like, and a Chinese hamster ovary cell CHO/DG44 cell, a rat myeloma YB2/0 cell and the host cells of the present invention described in the item 5 are preferred.

After introduction of the humanized antibody expression vector, a transformant capable of stably

producing the humanized antibody can be selected using a medium for animal cell culture comprising an agent such as G418 sulfate (hereinafter referred to as "G418"; manufactured by SIGMA) and the like in accordance with the method disclosed in Japanese Published Examined Patent Application No. 257891/90. Examples of the medium for animal cell culture include RPMI 1640 medium (manufactured by Nissui Pharmaceutical), GIT medium (manufactured by Nihon Pharmaceutical), EX-CELL 302 medium (manufactured by JRH), IMDM medium (manufactured by GIBCO BRL), Hybridoma-SFM medium (manufactured by GIBCO BRL) media obtained by adding various additives such as fetal bovine serum (hereinafter referred to as "FBS") to these media, and the like. The humanized antibody can be produced and accumulated in the culture supernatant by culturing the obtained transformant in a medium. The expression level and antigen binding activity of the humanized antibody in the culture supernatant can be measured by a method such as enzyme-linked immunosorbent assay [hereinafter referred to as "ELISA"; *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Chapter 14 (1998), *Monoclonal Antibodies: Principles and Practice*, Academic Press Limited (1996)] or the like. Also, the expression level of the humanized antibody by the transformant can be increased using a DHFR gene amplification system in accordance with the method disclosed in Japanese Published Examined Patent Application No. 257891/90.

The humanized antibody can be purified from a culture supernatant of the transformant using a protein A column [*Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Chapter 8 (1988), *Monoclonal Antibodies: Principles and Practice*, Academic Press Limited (1996)]. In addition, purification methods generally used for the purification of proteins can also be used. For example, the purification can be carried out through the combination

of a gel filtration, an ion exchange chromatography and an ultrafiltration. The molecular weight of the H chain, L chain or antibody molecule as a whole of the purified humanized antibody can be measured, e.g., by polyacrylamide gel electrophoresis [hereinafter referred to as "SDS-PAGE"; *Nature*, 227, 680 (1970)], Western blotting [*Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Chapter 12, (1988), *Monoclonal Antibodies: Principles and Practice*, Academic Press Limited (1996)] or the like.

Thus, methods for producing an antibody composition using an animal cell as the host have been described, but, as described above, the antibody composition can also be produced by a yeast, an insect cell, a plant cell, an animal individual or a plant individual by the same methods on the animal cell.

When a host cell has the ability to express an antibody molecule innately, the antibody composition of the present invention can be produced by preparing a cell expressing an antibody molecule using the method described in the item 1, culturing the cell and then purifying the antibody composition of interest from the resulting culture.

4. Activity evaluation of antibody composition

As the method for measuring the amount of the purified antibody composition, the activity to bind to an antibody and the effector function of the purified antibody composition, the known method described in *Monoclonal Antibodies*, *Antibody Engineering* and the like can be used.

As the examples, when the antibody composition is a humanized antibody, the binding activity with an antigen and the binding activity with an antigen-positive cultured cell line can be measured by methods such as ELISA, an immunofluorescent method [*Cancer Immunol. Immunother.*, 36, 373 (1993)] and the like. The cytotoxic activity against an antigen-positive cultured cell line can be evaluated by

Specifically, the compositional ratio can be calculated in accordance with a known method [*Agric. Biol. Chem.*, 55(1), 283-284 (1991)], by labeling an acid-hydrolyzed sample with a fluorescence with 2-aminopyridylation and then analyzing the composition by HPLC.

(2) Analysis of sugar chain structure

The sugar chain structure binding to an antibody molecule can be analyzed by the two dimensional sugar chain mapping method [*Anal. Biochem.*, 171, 73 (1988), *Biochemical Experimentation Methods 23 - Methods for Studying Glycoprotein Sugar Chains* (Japan Scientific Societies Press) edited by Reiko Takahashi (1989)]. The two dimensional sugar chain mapping method is a method for deducing a sugar chain structure by, e.g., plotting the retention time or elution position of a sugar chain by reverse phase chromatography as the X axis and the retention time or elution position of the sugar chain by normal phase chromatography as the Y axis, respectively, and comparing them with such results of known sugar chains.

Specifically, sugar chains are released from an antibody by subjecting the antibody to hydrazinolysis, and the released sugar chain is subjected to fluorescence labeling with 2-aminopyridine (hereinafter referred to as "PA") [*J. Biochem.*, 95, 197 (1984)], and then the sugar chains are separated from an excess PA-treating reagent by gel filtration, and subjected to reverse phase chromatography. Thereafter, each peak of the separated sugar chains are subjected to normal phase chromatography. The sugar chain structure can be deduced by plotting the results on a two dimensional sugar chain map and comparing them with the spots of a sugar chain standard (manufactured by Takara Shuzo) or a literature [*Anal. Biochem.*, 171, 73 (1988)].

The structure deduced by the two dimensional sugar chain mapping method can be confirmed by further carrying out mass spectrometry such as MALDI-TOF-MS of each sugar chain or the like.

6. Immunological determination method for discriminating sugar chain structure of antibody molecule

An antibody composition comprises an antibody molecule in which sugar chains binding to the Fc region of the antibody are different in structure. The antibody composition in which the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain is 20% or more among the total complex *N*-glycoside-linked sugar chains binding to the Fc region in the antibody composition reducing end has potent ADCC activity. The antibody composition can be identified by using the method for analyzing the sugar chain structure of an antibody molecule described in the item 6. Also, it can also be identified by an Immunological determination method using a lectin.

The sugar chain structure of an antibody molecule can be identified by the Immunological determination method using a lectin in accordance with the known Immunological determination method such as Western staining, IRA (radioimmunoassay), VIA (viroimmunoassay), EIA (enzymimmunoassay), FIA (fluoroimmunoassay), MIA (metalloimmunoassay) and the like described in *Monoclonal Antibodies: Principles and Applications*, Wiley-Liss, Inc. (1995); *Immunoassay*, 3rd Ed., Igakushoin (1987); *Enzyme Antibody Method*, Revised Edition, Gakusai Kikaku (1985); and the like.

A lectin which recognizes the sugar chain structure of an antibody molecule comprised in an antibody composition is labeled, and the labeled lectin is allowed to react with an antibody composition which is a sample.

It is preferable to use a lectin which specifically recognizes a sugar chain structure wherein fucose binds to the *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain. Examples include *Lens culinaris* lectin LCA (lentil agglutinin derived from *Lens culinaris*), pea lectin PSA (pea lectin derived from *Pisum sativum*), broad bean lectin VFA (agglutinin derived from *Vicia faba*) and *Aleuria aurantia* lectin AAL (lectin derived from *Aleuria aurantia*).

treated using balloon catheter at present, but circulatory organ diseases can be prevented and treated by inhibiting growth of arterial cells in restructure after treatment using an antibody having potent antibody-dependent cell-mediated cytotoxic activity.

Various diseases including viral and bacterial infections can be prevented and treated by inhibiting proliferation of cells infected with a virus or bacterium using an antibody having potent antibody-dependent cell-mediated cytotoxic activity.

Examples of an antibody which recognizes a tumor-related antigen, an antibody which recognizes an allergy- or inflammation-related antigen, an antibody which recognizes circulatory organ disease-related antigen and an antibody which recognizes a viral or bacterial infection-related antigen are described below.

Examples of the antibody which recognizes a tumor-related antigen include anti-GD2 antibody (Ohta et al., *Anticancer Res.*, 13, 331-336, 1993), anti-GD3 antibody (Ohta et al., *Cancer Immunol. Immunother.*, 36, 260-266, 1993), anti-GM2 antibody (Nakamura et al., *Cancer Res.*, 54, 1511-1516, 1994), anti-HER2 antibody (Carter et al., *Proc. Natl. Acad. Sci. USA*, 89, 4285-4289, 1992), anti-CD52 antibody (Carter et al., *Proc. Natl. Acad. Sci. USA*, 89, 4285-4289, 1992), anti-MAGE antibody (Jungbluth et al., *British J. Cancer*, 83, 493-497, 2000), anti-HM1.24 antibody (Ono et al., *Molecular Immunol.*, 36, 387-395, 1999), anti-parathyroid hormone-related protein (PTHrP) antibody (Ogata et al., *Cancer*, 88, 2909-2911, 2000), anti-basic fibroblast growth factor antibody and anti-FGF8 antibody (Matsuzaki et al., *Proc. Natl. Acad. Sci. USA*, 86, 9911-9915, 1989), anti-basic fibroblast growth factor receptor antibody and anti-FGF8 receptor antibody (Kuo et al., *J. Biol. Chem.*, 265, 16455-16463, 1990), anti-insulin-like growth factor antibody (Yao et al., *J. Neurosci. Res.*, 40, 647-659, 1995),

anti-insulin-like growth factor receptor antibody (Yao et al., *J. Neurosci. Res.*, 40, 647-659, 1995), anti-PMSA antibody (Murphy et al., *J. Urology*, 160, 2396-2401, 1998), anti-vascular endothelial cell growth factor antibody (Presta et al., *Cancer Res.*, 57, 4593-4599, 1997), anti-vascular endothelial cell growth factor receptor antibody (Kanno et al., *Oncogene*, 19, 2138-2146, 2000) and the like.

Examples of the antibody which recognizes an allergy- or inflammation-related antigen include anti-interleukin 6 antibody (Abrams et al., *Immunol. Rev.*, 127, 5-24, 1992), anti-interleukin 6 receptor antibody (Sato et al., *Molecular Immunol.*, 31, 371-381, 1994), anti-interleukin 5 antibody (Abrams et al., *Immunol. Rev.*, 127, 5-24, 1992), anti-interleukin 5 receptor antibody and anti-interleukin 4 antibody (Biord et al., *Cytokine*, 3, 562-567, 1991), anti-tumor necrosis factor antibody (Tempest et al., *Hybridoma*, 13, 183-190, 1994), anti-tumor necrosis factor receptor antibody (Amrani et al., *Molecular Pharmacol.*, 58, 237-245, 2000), anti-CCR4 antibody (Campbell et al., *Nature*, 400, 776-780, 1999), anti-chemokine antibody (Peri et al., *J. Immuno. Meth.*, 174, 249-257, 1994), anti-chemokine receptor antibody (Wu et al., *J. Exp. Med.*, 186, 1373-1381, 1997) and the like. Examples of the antibody which recognizes a circulatory organ disease-related antigen include anti-GpIIb/IIIa antibody (Co et al., *J. Immunol.*, 152, 2968-2976, 1994), anti-platelet-derived growth factor antibody (Ferns et al., *Science*, 253, 1129-1132, 1991), anti-platelet-derived growth factor receptor antibody (Shulman et al., *J. Biol. Chem.*, 272, 17400-17404, 1997) and anti-blood coagulation factor antibody (Peter et al., *Circulation*, 101, 1158-1164, 2000) and the like.

Examples of the antibody which recognizes a viral or bacterial infection-related antigen include anti-gp120 antibody (Tugarinov et al., *Structure*, 8, 385-395, 2000), anti-CD4 antibody (Schulze-Koops et al., *J. Rheumatology*,

polyethylene glycol, propylene glycol, etc.; oils, such as sesame oil, olive oil, soybean oil, etc.; antiseptics, such as p-hydroxybenzoic acid esters, etc.; flavors, such as strawberry flavor, peppermint, etc.; and the like.

Capsules, tablets, powders, granules and the like can be produced using, as additive, fillers, such as lactose, glucose, sucrose, mannitol, etc.; disintegrating agents, such as starch, sodium alginate, etc.; lubricants, such as magnesium stearate, talc, etc.; binders, such as polyvinyl alcohol, hydroxypropylcellulose, gelatin, etc.; surfactants, such as fatty acid ester, etc.; plasticizers, such as glycerine, etc.; and the like.

Examples of the pharmaceutical preparation suitable for parenteral administration include injections, suppositories, sprays and the like.

Injections may be prepared using a carrier, such as a salt solution, a glucose solution, a mixture of both thereof or the like. Also, powdered injections can be prepared by freeze-drying the antibody composition in the usual way and adding sodium chloride thereto.

Suppositories may be prepared using a carrier such as cacao butter, hydrogenated fat, carboxylic acid or the like.

Also, sprays may be prepared using the antibody composition as such or using a carrier which does not stimulate the buccal or airway mucous membrane of the patient and can facilitate absorption of the antibody composition by dispersing it as fine particles.

Examples of the carrier include lactose, glycerol and the like. Depending on the properties of the antibody composition and the carrier, it is possible to produce pharmaceutical preparations such as aerosols, dry powders and the like. In addition, the components exemplified as additives for oral preparations can also be added to the parenteral preparations.

Although the clinical dose or the frequency of administration varies depending on the objective therapeutic effect, administration method, treating period, age, body weight and the like, it is usually 10 µg/kg to 20 mg/kg per day and per adult.

Also, as the method for examining antitumor effect of the antibody composition against various tumor cells, *in vitro* tests include CDC activity measuring method, ADCC activity measuring method and the like, and *in vivo* tests include antitumor experiments using a tumor system in an experimental animal such as a mouse, etc. and the like.

CDC activity and ADCC activity measurements and antitumor experiments can be carried out in accordance with the methods described in *Cancer Immunology Immunotherapy*, 36, 373 (1993); *Cancer Research*, 54, 1511 (1994) and the like.

The present invention will be described below in detail based on Examples; however, Examples are only simple illustrations, and the scope of the present invention is not limited thereto.

Example 1

Production of anti-ganglioside GD3 human chimeric antibody:

1. Construction of tandem expression vector pChiLHGM4 for anti-ganglioside GD3 human chimeric antibody

A plasmid pChi641LGM40 was constructed by ligating a fragment of about 4.03 kb containing an L chain cDNA, obtained by digesting an L chain expression vector, pChi641LGM4 [*J. Immunol. Methods*, 167, 271 (1994)] for anti-ganglioside GD3 human chimeric antibody (hereinafter referred to as "anti-GD3 chimeric antibody") with restriction enzymes *MluI* (manufactured by Takara Shuzo) and *SalI* (manufactured by Takara Shuzo) with a fragment of about 3.40 kb containing a G418-resistant gene and a splicing signal, obtained by digesting an expression vector

PAGE107 [Cytotechnology, 3, 133 (1990)] for animal cell with restriction enzymes *Mlu*I (manufactured by Takara Shuzo) and *Sal*I (manufactured by Takara Shuzo) using DNA Ligation Kit (manufactured by Takara Shuzo), and then transforming *E. coli* HB101 (*Molecular Cloning*, Second Edition) with the ligated product.

Next, a fragment of about 5.68 kb containing an L chain cDNA, obtained by digesting the constructed plasmid pChi641LGM40 with a restriction enzyme *Cla*I (manufactured by Takara Shuzo), blunt-terminating it using DNA Blunting Kit (manufactured by Takara Shuzo) and further digesting it with *Mlu*I (manufactured by Takara Shuzo), was ligated with a fragment of about 8.40 kb containing an H chain cDNA, obtained by digesting an anti-GD3 chimeric antibody H chain expression vector, pChi641HGM4 [*J. Immunol. Methods*, 167, 271 (1994)] with a restriction enzyme, *Xho*I (manufactured by Takara Shuzo), blunt-terminating it using DNA Blunting Kit (manufactured by Takara Shuzo) and further digesting it with *Mlu*I (manufactured by Takara Shuzo), using DNA Ligation Kit (manufactured by Takara Shuzo), and then *E. coli* HB101 (*Molecular Cloning*, Second Edition) was transformed with the ligated product to thereby construct a tandem expression vector pChi641LHGM4 for anti-GD3 chimeric antibody.

2. Preparation of cells stably producing anti-GD3 chimeric antibody

Cells capable of stably producing an anti-GD3 chimeric antibody were prepared using the tandem expression vector pChi641LHGM4 for anti-GD3 chimeric antibody constructed in the item 1 of Example 1, as described below.

(1) Preparation of antibody-producing cell using rat myeloma YB2/0 cell

After introducing 5 μ g of the anti-GD3 chimeric antibody expression vector pChi641LHGM4 into 4×10^6 cells of rat myeloma YB2/0 [ATCC CRL-1662, J.V. Kilmarin et al., *J. Cell. Biol.*, 93, 576-582 (1982)] by electroporation [Cytotechnology, 3, 133 (1990)], the cells were suspended in 40 ml of RPMI1640-FBS(10) (RPMI1640 medium comprising 10% FBS (manufactured by GIBCO BRL)) and dispensed in 200 μ l/well into a 96 well culture plate (manufactured by Sumitomo Bakelite). After culturing them at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added to a concentration of 0.5 mg/ml, followed by culturing for 1 to 2 weeks. The culture supernatant was recovered from wells in which colonies of transformants showing G418 resistance were formed and growth of colonies was observed, and the antigen binding activity of the anti-GD3 chimeric antibody in the supernatant was measured by the ELISA shown in the item 3 of Example 1.

Regarding the transformants in wells in which production of the anti-GD3 chimeric antibody was observed in culture supernatants, in order to increase the amount of the antibody production using a DHFR gene amplification system, each of them was suspended in the RPMI1640-FBS(10) medium comprising 0.5 mg/ml G418 and 50 nM DHFR inhibitor, methotrexate (hereinafter referred to as "MTX"; manufactured by SIGMA) to give a density of 1 to 2×10^5 cells/ml, and the suspension was dispensed in 2 ml into wells of a 24 well plate (manufactured by Greiner). Transformants showing 50 nM MTX resistance were induced by culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator. The antigen binding activity of the anti-GD3 chimeric antibody in culture supernatants in wells in which growth of transformants was observed was measured by the ELISA shown in the item 3 of Example 1. Regarding the

transformants in wells in which production of the anti-GD3 chimeric antibody was observed in culture supernatants, the MTX concentration was increased to 100 nM and then to 200 nM, and transformants capable of growing in the RPMI1640-FBS(10) medium comprising 0.5 mg/ml G418 and 200 nM MTX and of producing the anti-GD3 chimeric antibody in a large amount were finally obtained by the same method as described above. Among the obtained transformants, suitable cell lines were selected and were made into a single cell (cloning) by limiting dilution twice. Also, using the method for determining the transcription product of an α -1,6-fucosyltransferase gene shown in Example 9, a cell line producing a relatively small amount of the transcription product was selected and used as a suitable cell line.

The obtained anti-GD3 chimeric antibody-producing transformed cell clone 7-9-51 has been deposited on April 5, 1999, as FERM BP-6691 in National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology (Higashi 1-1-3, Tsukuba, Ibaraki, Japan) (present name: International Patent Organism Depositary, National Institute of Advanced Industrial Science and Technology (AIST Tsukuba Central 6, 1-1, Higashi 1-Chome Tsukuba-shi, Ibaraki-ken, Japan)).

(2) Preparation of antibody-producing cell using CHO/DG44 cell

After introducing 4 μ g of the anti-GD3 chimeric antibody expression vector, pChi641LHGM4, into 1.6×10^6 cells of CHO/DG44 [G. Urlaub and L.A. Chasin, *Proc. Natl. Acad. Sci. USA*, **77**, 4216-4220 (1980)] by electroporation [Cytotechnology, **3**, 133 (1990)], the cells were suspended in 10 ml of IMDM-FBS(10) [IMDM medium comprising 10% FBS and 1x concentration of HT supplement (manufactured by GIBCO BRL)] and dispensed in 200 μ l/well into a 96 well

culture plate (manufactured by Iwaki Glass). After culturing them at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added to give a concentration of 0.5 mg/ml, followed by culturing for 1 to 2 weeks. The culture supernatant was recovered from wells in which colonies of transformants showing G418 resistance were formed and growth of colonies was observed, and the antigen binding activity of the anti-GD3 chimeric antibody in the supernatant was measured by the ELISA shown in the item 3 of Example 1.

Regarding the transformants in wells in which production of the anti-GD3 chimeric antibody was observed in culture supernatants, in order to increase the amount of the antibody production using a DHFR gene amplification system, each of them was suspended in an IMDM-dFBS(10) medium [IMDM medium comprising 10% dialyzed fetal bovine serum (hereinafter referred to as "dFBS"; manufactured by GIBCO BRL)] comprising 0.5 mg/ml G418 and 10 nM MTX to give a density of 1 to 2×10^5 cells/ml, and the suspension was dispensed in 0.5 ml into wells of a 24 well plate (manufactured by Iwaki Glass). Transformants showing 10 nM MTX resistance were induced by culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator. Regarding the transformants in wells in which their growth was observed, the MTX concentration was increased to 100 nM, and transformants capable of growing in the IMDM-dFBS(10) medium comprising 0.5 mg/ml G418 and 100 nM MTX and of producing the anti-GD3 chimeric antibody in a large amount were finally obtained by the same method as described above. Among the obtained transformants, suitable cell lines were selected and were made into a single cell (cloning) by limiting dilution twice. Also, using the method for determining the transcription product of an α -1,6-fucosyltransferase gene shown in Example 9, a cell line producing a relatively

in wells in which growth of transformants was observed was measured by the ELISA shown in the item 3 of Example 1. Regarding the transformants in wells in which production of the anti-GD3 chimeric antibody was observed in culture supernatants, the MTX concentration was increased to 100 nM and then to 200 nM, and transformants capable of growing in the EX-CELL302-dFBS(10) medium comprising 0.5 mg/ml G418 and 200 nM MTX and of producing the anti-GD3 chimeric antibody in a large amount was finally obtained by the same method as described above. Among the obtained transformants, elite cell lines were selected and were made into a single cell (cloning) by limiting dilution twice. Also, using the method for determining the transcription product of an α -1,6-fucosyltransferase gene shown in Example 9, a cell line producing a relatively small amount of the transcription product was selected and used as a suitable cell line.

3. Measurement of binding activity of antibody to GD3 (ELISA)

The binding activity of the antibody to GD3 was measured as described below.

In 2 ml of ethanol solution containing 10 μ g of dipalmitoylphosphatidylcholine (manufactured by SIGMA) and 5 μ g of cholesterol (manufactured by SIGMA), 4 nmol of GD3 was dissolved. Into each well of a 96 well plate for ELISA (manufactured by Greiner), 20 μ l of the solution (40 pmol/well in final concentration) was dispensed, followed by air-drying, 1% bovine serum albumin (hereinafter referred to as "BSA"; manufactured by SIGMA)-containing PBS (hereinafter referred to as "1% BSA-PBS") was dispensed in 100 μ l/well, and then the reaction was carried out at room temperature for 1 hour for blocking remaining active groups. After discarding 1% BSA-PBS, a culture supernatant of a transformant or a diluted solution of a human chimeric

antibody was dispensed in 50 μ l/well to carry out the reaction at room temperature for 1 hour. After the reaction, each well was washed with 0.05% Tween 20 (manufactured by Wako Pure Chemical Industries)-containing PBS (hereinafter referred to as "Tween-PBS"), a peroxidase-labeled goat anti-human IgG (H & L) antibody solution (manufactured by American Qualex) diluted 3,000 times with 1% BSA-PBS was dispensed in 50 μ l/well as a secondary antibody solution, and then the reaction was carried out at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, ABTS substrate solution [solution prepared by dissolving 0.55 g of 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) ammonium salt in 1 liter of 0.1 M citrate buffer (pH 4.2) and adding 1 μ l/ml of hydrogen peroxide to the solution just before use (hereinafter the same solution was used)] was dispensed in 50 μ l/well for color development, and then absorbance at 415 nm (hereinafter referred to as "OD415") was measured.

4. Purification of anti-GD3 chimeric antibody

(1) Culturing of antibody-producing cell derived from YB2/0 cell and purification of antibody

The anti-GD3 chimeric antibody-producing transformed cell clone obtained in the item 2(1) of Example 1 was suspended in the Hybridoma-SFM medium comprising 0.2% BSA, 200 nM MTX and 100 nM triiodothyronine (hereinafter referred to as "T3"; manufactured by SIGMA) to give a density of 3×10^5 cells/ml and cultured using a 2.0 liter capacity spinner bottle (manufactured by Iwaki Glass) under agitating at a rate of 50 rpm. After culturing them at 37°C for 10 days in a temperature-controlling room, the culture supernatant was recovered. The anti-GD3 chimeric antibody was purified from the culture supernatant using a Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's instructions. The

purified anti-GD3 chimeric antibody was named YB2/0-GD3 chimeric antibody.

(2) Culturing of antibody-producing cell derived from CHO/DG44 cell and purification of antibody

The anti-GD3 chimeric antibody-producing transformed cell clone obtained in the item 2(2) of Example 1 was suspended in the EX-CELL302 medium comprising 3 mM L-Gln, 0.5% fatty acid concentrated solution (hereinafter referred to as "CDLC"; manufactured by GIBCO BRL) and 0.3% Pluronic F68 (hereinafter referred to as "PF68"; manufactured by GIBCO BRL) to give a density of 1×10^6 cells/ml, and the suspension was dispensed in 50 ml into 175 mm² flasks (manufactured by Greiner). After culturing them at 37°C for 4 days in a 5% CO₂ incubator, the culture supernatant was recovered. The anti-GD3 chimeric antibody was purified from the culture supernatant using a Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's instructions. The purified anti-GD3 chimeric antibody was named CHO/DG44-GD3 chimeric antibody.

(3) Culturing of antibody-producing cell derived from NS0 cell and purification of antibody

The anti-GD3 chimeric antibody-producing transformed cell clone obtained in the item 2(3) of Example 1 was suspended in the EX-CELL302 medium comprising 2 mM L-Gln, 0.5 mg/ml G418, 200 nM MTX and 1% FBS, to give a density of 1×10^6 cells/ml, and the suspension was dispensed in 200 ml into 175 mm² flasks (manufactured by Greiner). After culturing them at 37°C for 4 days in a 5% CO₂ incubator, the culture supernatant was recovered. The anti-GD3 chimeric antibody was purified from the culture supernatant using a Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's

instructions. The purified anti-GD3 chimeric antibody was named NS0-GD3 chimeric antibody (302).

Also, the transformed cell clone was suspended in the GIT medium comprising 0.5 mg/ml G418 and 200 nM MTX to give a density of 3×10^5 cells/ml, and the suspension was dispensed in 200 ml into 175 mm² flasks (manufactured by Greiner). After culturing them at 37°C for 10 days in a 5% CO₂ incubator, the culture supernatant was recovered. The anti-GD3 chimeric antibody was purified from the culture supernatant using a Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's instructions. The purified anti-GD3 chimeric antibody was named NS0-GD3 chimeric antibody (GIT).

(4) Culturing of antibody-producing cell derived from SP2/0 cell and purification of antibody

The anti-GD3 chimeric antibody-producing transformed cell clone (KM-871 (FERM BP-3512)) described in Japanese Published Unexamined Patent Application No. 304989/93 (EP 533199) was suspended in the GIT medium comprising 0.5 mg/ml G418 and 200 nM MTX to give a density of 3×10^5 cells/ml, and the suspension was dispensed in 200 ml into 175 mm² flasks (manufactured by Greiner). After culturing them at 37°C for 8 days in a 5% CO₂ incubator, the culture supernatant was recovered. The anti-GD3 chimeric antibody was purified from the culture supernatant using a Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's instructions. The purified anti-GD3 chimeric antibody was named SP2/0-GD3 chimeric antibody.

5. Analysis of purified anti-GD3 chimeric antibody

In accordance with a known method [Nature, 227, 680 (1970)], 4 µg of each of the five kinds of the anti-GD3 chimeric antibodies produced by and purified from

binding activity measured by changing the concentration of the anti-GD3 chimeric antibody to be added. As shown in Fig. 2, the five kinds of the anti-GD3 chimeric antibodies showed almost the same binding activity to GD3. The result shows that antigen binding activities of these antibodies are constant independently of the antibody-producing animal cells and their culturing methods. Also, it was suggested from the comparison of the NS0-GD3 chimeric antibody (302) with the NS0-GD3 chimeric antibody (GIT) that the antigen binding activities are constant independently of the media used in the culturing.

2. *In vitro* cytotoxic activity (ADCC activity) of anti-GD3 chimeric antibody

In order to evaluate *in vitro* cytotoxic activity of the five kinds of the purified anti-GD3 chimeric antibodies obtained in the item 4 of Example 1, the ADCC activity was measured in accordance with the following method.

(1) Preparation of target cell solution

A human melanoma cultured cell line G-361 (ATCC CRL 1424) was cultured using the RPMI1640-FBS(10) medium to prepare 1×10^6 cells, and the cells were radioisotope-labeled by reacting them with 3.7 MBq equivalents of a radioactive substance $\text{Na}_2^{51}\text{CrO}_4$ at 37°C for 1 hour. After the reaction, the cells were washed three times through their suspension in the RPMI1640-FBS(10) medium and centrifugation, re-suspended in the medium and then incubated at 4°C for 30 minutes in ice for spontaneous dissolution of the radioactive substance. After centrifugation, the precipitate was adjusted to 2×10^5 cells/ml by adding 5 ml of the RPMI1640-FBS(10) medium and used as the target cell solution.

(2) Preparation of effector cell solution

From a healthy person, 50 ml of venous blood was collected, and gently mixed with 0.5 ml of heparin sodium (manufactured by Takeda Pharmaceutical). The mixture was centrifuged to isolate a mononuclear cell layer using Lymphoprep (manufactured by Nycomed Pharma AS) in accordance with the manufacture's instructions. After washing with the RPMI1640-FBS(10) medium by centrifugation three times, the resulting precipitate was re-suspended to give a density of 2×10^6 cells/ml using the medium and used as the effector cell solution.

(3) Measurement of ADCC activity

Into each well of a 96 well U-shaped bottom plate (manufactured by Falcon), 50 μ l of the target cell solution prepared in the above (1) (1×10^4 cells/well) was dispensed. Next, 100 μ l of the effector cell solution prepared in the above (2) was added thereto (2×10^5 cells/well, the ratio of effector cells to target cells becomes 20:1). Subsequently, each of the anti-GD3 chimeric antibodies was added to give a final concentration from 0.0025 to 2.5 μ g/ml, followed by reaction at 37°C for 4 hours. After the reaction, the plate was centrifuged, and the amount of ^{51}Cr in the supernatant was measured using a γ -counter. The amount of spontaneously released ^{51}Cr was calculated by the same operation using only the medium instead of the effector cell solution and the antibody solution, and measuring the amount of ^{51}Cr in the supernatant. The amount of total released ^{51}Cr was calculated by the same operation using only the medium instead of the antibody solution and adding 1 N hydrochloric acid instead of the effector cell solution, and measuring the amount of ^{51}Cr in the supernatant. The ADCC activity was calculated from the following equation (II):

$$\text{ADCC activity (\%)} = \frac{{}^{51}\text{Cr in sample supernatant} - \text{spontaneously released } {}^{51}\text{Cr}}{\text{total released } {}^{51}\text{Cr} - \text{spontaneously released } {}^{51}\text{Cr}} \times 100 \text{ (II)}$$

The results are shown in Fig. 3. As shown in Fig. 3, among the five kinds of the anti-GD3 chimeric antibodies, the YB2/0-GD3 chimeric antibody showed the most potent ADCC activity, followed by the SP2/0-GD3 chimeric antibody, NS0-GD3 chimeric antibody and CHO-GD3 chimeric antibody in that order. No difference in the ADCC activity was found between the NS0-GD3 chimeric antibody (302) and NS0-GD3 chimeric antibody (GIT) prepared using different media in the culturing. The above results show that the ADCC activity of antibodies greatly varies depending on the kind of the animal cells to be used in their production. As its mechanism, since their antigen binding activities were identical, it was considered that it is caused by a difference in the structure binding to the antibody Fc region.

Example 3

Preparation of anti-human interleukin 5 receptor α chain human CDR-grafted antibody:

1. Preparation of cell stably producing anti-human interleukin 5 receptor α chain human CDR-grafted antibody
- (1) Preparation of antibody-producing cell using rat myeloma YB2/0 cell

Using the anti-human interleukin 5 receptor α chain human CDR-grafted antibody (hereinafter referred to as "anti-hIL-5R α CDR-grafted antibody") expression vector, pKANTEX1259HV3LV0, described in WO 97/10354, cells capable of stably producing anti-hIL-5R α CDR-grafted antibody were prepared as described below.

After introducing 5 μ g of the anti-hIL-5R α CDR-grafted antibody expression vector pKANTEX1259HV3LV0 into 4×10^6 cells of rat myeloma YB2/0 by electroporation

[Cytotechnology, 3, 133 (1990)], the cells were suspended in 40 ml of RPMI1640-FBS(10) and dispensed in 200 μ l/well into a 96 well culture plate (manufactured by Sumitomo Bakelite). After culturing them at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added to give a concentration of 0.5 mg/ml, followed by culturing for 1 to 2 weeks. The culture supernatant was recovered from wells in which colonies of transformants showing G418 resistance were formed and growth of colonies was observed, and the antigen binding activity of the anti-hIL-5R α CDR-grafted antibody in the supernatant was measured by the ELISA shown in the item 2 of Example 3.

Regarding the transformants in wells in which production of the anti-hIL-5R α CDR-grafted antibody was observed in culture supernatants, in order to increase amount of the antibody production using a DHFR gene amplification system, each of the them was suspended in the RPMI1640-FBS(10) medium comprising 0.5 mg/ml G418 and 50 nM MTX to give a density of 1 to 2×10^5 cells/ml, and the suspension was dispensed in 2 ml into wells of a 24 well plate (manufactured by Greiner). Transformants showing 50 nM MTX resistance were induced by culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator. The antigen binding activity of the anti-hIL-5R α CDR-grafted antibody in culture supernatants in wells in which growth of transformants was observed was measured by the ELISA shown in the item 2 of Example 3. Regarding the transformants in wells in which production of the anti-hIL-5R α CDR-grafted antibody was observed in culture supernatants, the MTX concentration was increased to 100 nM and then to 200 nM, and transformants capable of growing in the RPMI1640-FBS(10) medium comprising 0.5 mg/ml G418 and 200 nM MTX and of producing the anti-hIL-5R α CDR-grafted antibody in a large amount were finally obtained in the same manner as described above. Among the obtained transformants, elite

cell lines were selected and were made into a single cell (cloning) by limiting dilution twice. Also, using the method for determining the transcription product of an α -1,6-fucosyltransferase gene shown in Example 9, a cell line producing a relatively small amount of the transcription product was selected and used as a suitable cell line. The obtained anti-hIL-5R α CDR-grafted antibody-producing transformed cell clone No. 3 has been deposited on April 5, 1999, as FERM BP-6690 in National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology (Higashi 1-1-3, Tsukuba, Ibaraki, Japan) (present name: International Patent Organism Depositary, National Institute of Advanced Industrial Science and Technology (AIST Tsukuba Central 6, 1-1, Higashi 1-Chome Tsukuba-shi, Ibaraki-ken, Japan)).

(2) Preparation of antibody-producing cell using CHO/dhfr⁻ cell

After introducing 4 μ g of the anti-hIL-5R α CDR-grafted antibody expression vector pKANTEX1259HV3LV0 described in WO 97/10354 into 1.6×10^6 cells of CHO/dhfr⁻ by electroporation [Cytotechnology, 3, 133 (1990)], the cells were suspended in 10 ml of IMDM-FBS(10) and dispensed in 200 μ l/well into a 96 well culture plate (manufactured by Iwaki Glass). After culturing them at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added to give a concentration of 0.5 mg/ml, followed by culturing for 1 to 2 weeks. The culture supernatant was recovered from respective well in which colonies of transformants showing G418 resistance were formed and growth of colonies was observed, and the antigen binding activity of the anti-hIL-5R α CDR-grafted antibody in the supernatant was measured by the ELISA shown in the item 2 of Example 3.

Regarding the transformants in wells in which production of the anti-hIL-5R α CDR-grafted antibody was

observed in culture supernatants, in order to increase amount of the antibody production using a DHFR gene amplification system, each of the transformants was suspended in an IMDM-dFBS(10) medium comprising 0.5 mg/ml G418 and 10 nM MTX to give a density of 1 to 2×10^5 cells/ml, and the suspension was dispensed in 0.5 ml into wells of a 24 well plate (manufactured by Iwaki Glass). Transformants showing 10 nM MTX resistance were induced by culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator. Regarding the transformants in wells in which their growth was observed, the MTX concentration was increased to 100 nM and then to 500 nM, and transformants capable of growing in the IMDM-dFBS(10) medium comprising 0.5 mg/ml G418 and 500 nM MTX and of producing the anti-hIL-5R α CDR-grafted antibody in a large amount were finally obtained in the same manner as described above. Among the obtained transformants, elite cell lines were selected and were made into a single cell (cloning) by limiting dilution twice. Also, using the method for determining the transcription product of an α -1,6-fucosyltransferase gene shown in Example 9, a cell line producing a relatively small amount of the transcription product was selected and used as a suitable cell line.

(3) Production of antibody-producing cell using mouse myeloma NS0 cell

An anti-hIL-5R α CDR-grafted antibody expression vector was prepared in accordance with the method of Yarranton et al. [BIO/TECHNOLOGY, 10, 169 (1992)] and using the antibody H chain cDNA and L chain cDNA on the anti-hIL-5R α CDR-grafted antibody expression vector pKANTEX1259HV3LV0 described in WO 97/10354, and NS0 cell was transformed to obtain transformants capable of producing the anti-hIL-5R α CDR-grafted antibody in a large amount. Among the obtained transformants, elite cell lines

were selected and were made into a single cell (cloning) by limiting dilution twice. Also, using the method for determining the transcription production of an α -1,6-fucosyltransferase gene shown in Example 9, a cell line producing a relatively small amount of the transcription product was selected and used as a suitable cell line.

2. Measurement of binding activity of antibody to hIL-5R α (ELISA)

The binding activity of the antibody to hIL-5R α was measured as described below.

A solution was prepared by diluting the anti-hIL-5R α mouse antibody KM1257 described in WO 97/10354 with PBS to give a concentration of 10 μ g/ml, and 50 μ l of the resulting solution was dispensed into each well of a 96 well plate for ELISA (manufactured by Greiner), followed by reaction at 4°C for 20 hours. After the reaction, 1% BSA-PBS was dispensed in 100 μ l/well, and then the reaction was carried out at room temperature for 1 hour to block remaining active groups. After discarding 1% BSA-PBS, a solution prepared by diluting the soluble hIL-5R α described in WO 97/10354 with 1% BSA-PBS to give a concentration of 0.5 μ g/ml was dispensed in 50 μ l/well, followed by reaction at 4°C for 20 hours. After the reaction, each well was washed with Tween-PBS, culture supernatants of transformants or diluted solutions of a purified human CDR-grafted antibodies were dispensed in 50 μ g/well to carry out the reaction at room temperature for 2 hours. After the reaction, each well was washed with Tween-PBS, a peroxidase-labeled goat anti-human IgG (H & L) antibody solution (manufactured by American Qualex) diluted 3,000 times with 1% BSA-PBS was dispensed in 50 μ l/well as a secondary antibody solution, followed by reaction at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, the ABTS substrate solution was

(3) Culturing of antibody-producing cell derived from NS0 cell and purification of antibody

The anti-hIL-5R α CDR-grafted antibody-producing transformed cell clone obtained in the item 1(3) of Example 3 was cultured in accordance with the method of Yarranton et al. [BIO/TECHNOLOGY, 10, 169 (1992)] and then a culture supernatant was recovered. The anti-hIL-5R α CDR-grafted antibody was purified from the culture supernatant using ion exchange chromatography and the gel filtration method. The purified anti-hIL-5R α CDR-grafted antibody was named NS0-hIL-5R CDR antibody.

4. Analysis of purified anti-hIL-5R α CDR-grafted antibodies

In accordance with a known method [Nature, 227, 680 (1970)], 4 μ g of each of the three kinds of the anti-hIL-5R α CDR-grafted antibodies produced by and purified from each animal cells, obtained in the item 3 of Example 3, was subjected to SDS-PAGE to analyze the molecular weight and purification degree. The results are shown in Fig. 4. As shown in Fig. 4, a single band of about 150 Kd in molecular weight was found under non-reducing conditions, and two bands of about 50 Kd and about 25 Kd under reducing conditions, in each of the purified anti-hIL-5R α CDR-grafted antibodies. The molecular weights almost coincided with the molecular weights deduced from the cDNA nucleotide sequences of H chain and L chain of the antibody (H chain: about 49 Kd, L chain: about 23 Kd, whole molecule: about 144 Kd), and also coincided with the reports stating that the IgG antibody has a molecular weight of about 150 Kd under non-reducing conditions and is degraded into H chains having a molecular weight of about 50 Kd and L chains having a molecular weight of about 25 Kd under reducing conditions due to cutting of the S-S bond in the molecule [Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Chapter 14 (1998); Monoclonal Antibodies:

Principles and Practice, Academic Press Limited (1996)], so that it was confirmed that each anti-hIL-5R α CDR-grafted antibody was expressed and purified as an antibody molecule having the true structure.

Example 4

Activity evaluation of anti-hIL-5R α CDR-grafted antibody:

1. Binding activity of anti-hIL-5R α CDR-grafted antibody to hIL-5R α (ELISA)

The activity of the three kinds of the purified anti-hIL-5R α CDR-grafted antibodies obtained in the item 3 of Example 3 to bind to hIL-5R α was measured by the ELISA shown in the item 2 of Example 3. Fig. 5 shows a result of the examination of the binding activity measured by changing concentration of the anti-hIL-5R α CDR-grafted antibody to be added. As shown in Fig. 5, the three kinds of the anti-hIL-5R α CDR-grafted antibodies showed almost the same binding activity to hIL-5R α . The result shows that the antigen binding activities of these antibodies are constant independently of the antibody-producing animal cells and their culturing methods, similar to the result of the item 1 of Example 2.

2. *In vitro* cytotoxic activity (ADCC activity) of anti-hIL-5R α CDR-grafted antibody

In order to evaluate *in vitro* cytotoxic activity of the three kinds of the purified anti-hIL-5R α CDR-grafted antibodies obtained in the item 3 of Example 3, the ADCC activity was measured in accordance with the following method.

(1) Preparation of target cell solution

A mouse T cell line CTLL-2(h5R) expressing the hIL-5R α chain and β chain described in WO 97/10354 was cultured using the RPMI1640-FBS(10) medium to give a density of

(2) Preparation of effector cell solution

(3) Measurement of ADCC activity

- 136 -

amount of ^{51}Cr in the supernatant was measured using a γ -counter. The amount of spontaneously released ^{51}Cr was calculated by the same operation using only the medium instead of the effector cell solution and the antibody solution, and measuring the amount of ^{51}Cr in the supernatant. The amount of total released ^{51}Cr was calculated by the same operation using only the medium instead of the antibody solution and adding 1 N hydrochloric acid instead of the effector cell solution, and measuring the amount of ^{51}Cr in the supernatant.

The ADCC activity was calculated from the above equation (II).

The results are shown in Fig. 6. As shown in Fig. 6, among the three kinds of the anti-hIL-5R α CDR-grafted antibodies, the YB2/0-hIL-5R CDR antibody showed the most potent ADCC activity, followed by the CHO/d-hIL-5R CDR antibody and the NS0-hIL-5R CDR antibody in this order. Similar to the result of the item 2 of Example 2, the above results show that the ADCC activity of antibodies greatly varies depending on the animal cells to be used in their production. In addition, since the antibodies produced by the YB2/0 cell showed the most potent ADCC activity in both cases of the two kinds of the humanized antibodies, it was revealed that an antibody having potent ADCC activity can be produced by using the YB2/0 cell.

3. *In vivo* activity evaluation of anti-hIL-5R α CDR-grafted antibody

In order to evaluate *in vivo* activity of the three kinds of the purified anti-hIL-5R α CDR-grafted antibodies obtained in the item 3 of Example 3, the inhibition activity in an hIL-5-induced eosinophilia increasing model of *Macaca fascicularis* was examined in accordance with the following method.

The hIL-5 (preparation method is described in WO 97/10354) was administered to *Macaca fascicularis* under the dorsal skin at a dose of 1 µg/kg, starting on the first day and once a day for a total of 14 times. Each anti-hIL-5Rα CDR-grafted antibody was intravenously administered at a dose of 0.3 mg/kg one hour before the hIL-5 administration on the day zero. An antibody-non-added group was used as the control. In the antibody-administered groups, three animals of *Macaca fascicularis* were used in each group (No. 301, No. 302, No. 303, No. 401, No. 402, No. 403, No. 501, No. 502 and No. 503), and two animals (No. 101 and No. 102) were used in the antibody-non-added group. Starting 7 days before commencement of the administration and until 42 days after the administration, about 1 ml of blood was periodically collected from a saphena or a femoral vein, and the number of eosinophils in 1 µl of peripheral blood was measured. The results are shown in Fig. 7. As shown in Fig. 7, increase in the blood eosinophil was completely inhibited in the group to which the YB2/0-hIL-5R CDR antibody was administered. On the other hand, complete inhibition activity was found in one animal in the group to which the CHO/d-hIL-5R CDR antibody was administered, but the inhibition activity was not sufficient in two animals. In the group to which NS0-hIL-5R CDR antibody was administered, complete inhibition activity was not found and its effect was not sufficient.

The above results show that the *in vivo* activity of antibodies greatly varies depending on the animal cells to be used in their production. In addition, since a positive correlation was found between the degree of the *in vivo* activity of the anti-hIL-5Rα CDR-grafted antibody and the degree of its ADCC activity described in the item 2 of Example 4, it was indicated that the degree of ADCC activity is remarkably important for its activity expression.

Based on the above results, it is expected that an antibody having potent ADCC activity is useful also in the clinical field for various diseases in human.

Example 5

Analysis of sugar chain which enhances ADCC activity:

1. Preparation of 2-aminopyridine-labeled sugar chain (PA-treated sugar chain)

The humanized antibody of the present invention was acid-hydrolyzed with hydrochloric acid to remove sialic acid. After hydrochloric acid was completely removed, the sugar chain was cleaved from the protein by hydrazinolysis [*Method of Enzymology*, 83, 263 (1982)]. Hydrazine was removed, and *N*-acetylation was carried out by adding an aqueous ammonium acetate solution and acetic anhydride. After lyophilizing, fluorescence labeling with 2-aminopyridine was carried out [*J. Biochem.*, 95, 197 (1984)]. The fluorescence-labeled sugar chain (PA-treated sugar chain) was separated from impurity using Surperdex Peptide HR 10/30 Column (manufactured by Pharmacia). The sugar chain fraction was dried using a centrifugal concentrator and used as a purified PA-treated sugar chain.

2. Reverse phase HPLC analysis of PA-treated sugar chain of purified anti-hIL-5R α CDR-grafted antibody

According to the method in the item 1 of Example 5, various anti-hIL-5R α CDR-grafted antibodies produced in Example 3 were subjected to PA-treated sugar chain treatment, and reverse phase HPLC analysis was carried out by CLC-ODS column (manufactured by Shimadzu). An excess amount of α -L-fucosidase (derived from bovine kidney, manufactured by SIGMA) was added to the PA-treated sugar chain for digestion (37°C, 15 hours), and then the products were analyzed by reverse phase HPLC (Fig. 8). It was confirmed that the asparagine-linked sugar chain is eluted

for 30 minutes to 80 minute using PA-treated sugar chain standards manufactured by Takara Shuzo. The ratio of sugar chains whose reverse phase HPLC elution positions were shifted (sugar chains eluted for 48 minutes to 78 minutes) by the α -L-fucosidase digestion was calculated. The results are shown in Table 1.

Table 1

Antibody-producing cell	α -1,6-Fucose-linked sugar chain (%)
YB2/0	47
NS0	73

About 47% of the anti-hIL-5R CDR-grafted antibody produced by the YB2/0 cell and about 73% of the anti-hIL-5R CDR-grafted antibody produced by the NS0 cell were sugar chains in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain (hereinafter referred to as "sugar chain having α -1,6-fucose"). Thus, the ratio of sugar chains in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain (hereinafter referred to as " α -1,6-fucose-free sugar chain") is higher in the antibody produced by the YB2/0 cell than in the antibody produced by the NS0 cell.

3. Analysis of monosaccharide composition of purified anti-hIL-5R α CDR-grafted antibody

Sugar chains of anti-hIL-5R α CDR-grafted antibodies produced by the YB2/0 cell, NS0 cell and CHO/d cell were hydrolyzed into monosaccharides by acid hydrolysis with trifluoroacetic acid, and monosaccharide composition analysis was carried out using BioLC (manufactured by Dionex).

Among *N*-glycoside-linked sugar chains, there are 3 mannose units in one sugar chain in the complex type *N*-glycoside-linked sugar chain. A relative ratio of each monosaccharide obtained by calculating the number of mannose as 3 is shown in Table 2.

Table 2

Antibody-producing cell	Fuc	GlcNAc	Gal	Man	ADCC activity (%)*
YB2/0	0.60	4.98	0.30	3.00	42.27
NS0	1.06	3.94	0.66	3.00	16.22
CHO/dhFr ⁻	0.85	3.59	0.49	3.00	25.73
CHO/dhFr ⁻	0.91	3.80	0.27	3.00	25.73

*: Antibody concentration: 0.01 µg/ml

Since the relative ratios of fucose were in an order of YB2/0 < CHO/d < NS0, the sugar chain produced in the antibody produced by YB2/0 cell showed the lowest fucose content as also shown in the present results.

4. Sugar chain analysis of antibody produced by CHO/dhfr⁻ cell

PA-treated sugar chains were prepared from purified anti-hIL-5Rα CDR-grafted antibody produced by CHO/dhfr⁻ cell, and reverse phase HPLC analysis was carried out using CLC-ODS column (manufactured by Shimadzu) (Fig. 9). In Fig. 9, an elution time from 35 to 45 minutes corresponded to sugar chains having no fucose and an elution time from 45 to 60 minutes corresponded to sugar chains having fucose. Similar to the case of the antibody produced by mouse myeloma NS0 cell, the anti-hIL-5Rα CDR-grafted antibody produced by CHO/dhfr⁻ cell had less fucose-free sugar chain content than the antibody produced by rat myeloma YB2/0 cell.

Example 6

Separation of potent ADCC activity antibody:

The anti-hIL-5R α CDR-grafted antibody produced by rat myeloma YB2/0 cell was separated using a lectin column which binds to sugar chains having fucose. HPLC was carried out using LC-6A manufactured by Shimadzu at a flow rate of 1 ml/min and at room temperature as the column temperature. After equilibration with 50 mM Tris-sulfate buffer (pH 7.3), the purified anti-hIL-5R α CDR-grafted antibody was injected and then eluted by a linear density gradient (60 minutes) of 0.2 M α -methylmannoside (manufactured by Nakalai Tesque). The anti-hIL-5R α CDR-grafted antibody was separated into non-adsorbed fraction and adsorbed fraction. When the non-adsorbed fraction and a part of the adsorbed fraction were sampled and their binding activity to hIL-5R α was measured, they showed similar binding activity (Fig. 10A). When the ADCC activity was measured, the non-adsorbed fraction showed potent ADCC activity (100 to 1000 folds) than that of the part of adsorbed fraction (Fig. 10B). In addition, PA-treated sugar chains were prepared from the non-adsorbed fraction and a part of the adsorbed fraction, and reverse HPLC analysis was carried out using CLC-ODS column (manufactured by Shimadzu) (Fig. 11). In the non-adsorbed fraction, an antibody binding to fucose-free sugar chains was mainly present, and in the part of adsorbed fraction, an antibody binding to sugar chains having fucose was mainly present.

Example 7

Activity evaluation of anti-GD3 chimeric antibody having different ratio of α -1,6-fucose-free sugar chain:

1. Preparation of anti-GD3 chimeric antibodies having different ratio of α -1,6-fucose-free sugar chain

In accordance with the method described in the item 2(1) of Example 1, transformed clones derived from YB2/0 cell capable of producing an anti-GD3 chimeric antibody was obtained. Antibodies were prepared from the transformed clones derived from YB2/0 cell and named lot 1, lot 2 and lot 3. Each sugar chain which is bound to the anti-GD3 chimeric antibodies of lot 1, lot 2 and lot 3 was analyzed in accordance with the method of Example 11(6), and it was found that the ratios of α -1,6-fucose-free sugar chains were 50%, 45% and 29%, respectively. Herein, these samples are referred to as anti-GD3 chimeric antibody (50%), anti-GD3 chimeric antibody (45%) and anti-GD3 chimeric antibody (29%).

Also, sugar chains of the anti-GD3 chimeric antibody derived from the CHO/DG44 cell prepared in the item 2(2) of Example 1 were analyzed in accordance with the method of Example 11(6), and it was found that the ratio of α -1,6-fucose-free sugar chains was 7%. Herein, the sample is referred to as anti-GD3 chimeric antibody (7%).

The anti-GD3 chimeric antibody (45%) and anti-GD3 chimeric antibody (7%) were mixed at a ratio of anti-GD3 chimeric antibody (45%) : anti-GD3 chimeric antibody (7%) = 5:3 or 1:7. Sugar chains of the samples were analyzed in accordance with the method of Example 10(6), and it was found that samples having the ratio of α -1,6-fucose-free sugar chains of 24% and 13% (calculated value) were prepared. Herein, they are referred to as anti-GD3 chimeric antibody (24%) and anti-GD3 chimeric antibody (13%).

Results of the sugar chain analysis of each of the samples are shown in Fig. 12. The ratio of α -1,6-fucose-free sugar chains was shown as an average value of the result of two sugar chain analyses.

2. Evaluation of binding activity to GD3 (ELISA)

The binding activities of the six kinds of the anti-GD3 chimeric antibodies having a different ratio of α -1,6-fucose-free sugar chains prepared in the item 1 of Example 7 against GD3 (manufactured by Snow Brand Milk Products) were measured by the ELISA shown in the item 3 of Example 1. As a result, all of the six kinds of the anti-GD3 chimeric antibodies showed almost the same GD3-binding activity as shown in Fig. 13, and it was found that the ratio of the α -1,6-fucose-free sugar chains does not have influence on the antigen binding activity of the antibody.

3. Evaluation of ADCC activity on human melanoma cell line

The ADCC activity of anti-GD3 chimeric antibodies on a human melanoma cell line G-361 (ATCC CRL 1424) was measured as follows.

(1) Preparation of target cell suspension

1×10^6 cells of a human melanoma cell line G-361 were prepared, a 3.7 MBq equivalent of a radioactive substance $\text{Na}_2^{51}\text{CrO}_4$ was added thereto and the mixture was allowed to react at 37°C for 1 hour to label the cells with the radioisotope. After the reaction, the cells were washed three times by a procedure of their suspension in a medium and subsequent centrifugation, re-suspended in the medium and then incubated at 4°C for 30 minutes in ice to effect spontaneous dissociation of the radioactive substance. After centrifugation, the cells were adjusted to 2×10^5 cells/ml by adding 5 ml of the medium and used as a target cell suspension.

(2) Preparation of human effector cell suspension

A 50 ml portion of peripheral blood was collected from a healthy person and gently mixed with 0.5 ml of heparin sodium (manufactured by Shimizu Pharmaceutical).

Using Lymphoprep (manufactured by AXIS SHIELD), this was centrifuged (800 g, 20 minutes) in accordance with the manufacture's instructions to separate a mononuclear cell layer. The cells were washed by centrifuging (1,200 rpm, 5 minutes) three times using a medium and then re-suspended in the medium to give a density of 2×10^6 cells/ml and used as a human effector cell suspension.

(3) Measurement of ADCC activity

The target cell suspension prepared in the (1) was dispensed in 50 μ l (1×10^4 cells/well) into each well of a 96 well U-bottom plate (manufactured by Falcon). Next, 100 μ l of the human effector cell suspension prepared in the (2) was added thereto (2×10^5 cells/well, ratio of the human effector cells to the target cells was 20 : 1). Each of the anti-GD3 chimeric antibodies was added thereto to give a final concentration of 0.0005 to 5 μ g/ml, followed by reaction at 37°C for 4 hours. After the reaction, the plate was centrifuged and the amount of ^{51}Cr in the supernatant was measured using a γ -counter. An amount of the spontaneously dissociated ^{51}Cr was calculated by carrying out the same procedure using the medium alone instead of the human effector cell suspension and antibody solution, and measuring the amount of ^{51}Cr in the supernatant. An amount of the total dissociated ^{51}Cr was calculated by carrying out the same procedure using the medium alone instead of the antibody solution and a 1 mol/l hydrochloric acid solution instead of the human effector cell suspension, and measuring the amount of ^{51}Cr in the supernatant. The cytotoxic activity (%) was calculated using equation (II).

Figs. 14 and 15 show results of the measurement of ADCC activity of the six kinds of the anti-GD3 chimeric antibodies having a different ratio of α -1,6-fucose-free sugar chains at various concentrations (0.0005 to 5 μ g/ml)

using effector cells of two healthy donors (A and B), respectively. As shown in Figs. 14 and 15, ADCC activity of the anti-GD3 chimeric antibodies showed a tendency to increase in proportion to the ratio of α -1,6-fucose-free sugar chains at each antibody concentration. The ADCC activity decreases when the antibody concentration is low. At an antibody concentration of 0.05 μ g/ml, the antibody in which the ratio of α -1,6-fucose-free sugar chains is 24%, 29%, 45% or 50% showed almost the same potent ADCC activity but the ADCC activity was low in the antibody (13%) or (7%) in which the ratio of α -1,6-fucose-free sugar chains is less than 20%. These results were the same when the effector cell donor was changed.

Example 8

Activity evaluation of anti-CCR4 chimeric antibody having different ratio of α -1,6-fucose-free sugar chain:

1. Production of cells stably producing anti-CCR4 chimeric antibody

Cells which capable of stably producing an anti-CCR4 chimeric antibody were prepared as follows using a tandem type expression vector pKANTEX2160 for an anti-CCR4 chimeric antibody described in WO 01/64754.

(1) Preparation of antibody-producing cell using rat myeloma YB2/0 cell

After introducing 10 μ g of the anti-CCR4 chimeric antibody expression vector pKANTEX2160 into 4×10^6 cells of rat myeloma YB2/0 cell (ATCC CRL 1662) by electroporation [Cytotechnology, 3, 133 (1990)], the cells were suspended in 40 ml of Hybridoma-SFM-FBS(5) [Hybridoma-SFM medium (manufactured by Invitrogen) comprising 5% FBS (manufactured by PAA Laboratories)] and dispensed in 200 μ l/well into a 96 well culture plate (manufactured by Sumitomo Bakelite). After culturing them at 37°C for 24

hours in a 5% CO₂ incubator, G418 was added to give a concentration of 1 mg/ml, followed by culturing for 1 to 2 weeks. Culture supernatant was recovered from wells in which growth of transformants showing G418 resistance was observed by the formation of colonies, and antigen binding activity of the anti-CCR4 chimeric antibody in the supernatant was measured by the ELISA described in the item 2 of Example 8.

Regarding the transformants in wells in which production of the anti-CCR4 chimeric antibody was observed in culture supernatants, in order to increase an amount of the antibody production using a DHFR gene amplification system, each of them was suspended in the Hybridoma-SFM-FBS(5) medium comprising 1 mg/ml G418 and 50 nM DHFR inhibitor MTX (manufactured by SIGMA) to give a density of 1 to 2 x 10⁵ cells/ml, and the suspension was dispensed in 1 ml portions into wells of a 24 well plate (manufactured by Greiner). After culturing them at 37°C for 1 to 2 weeks in a 5% CO₂ incubator, transformants showing 50 nM MTX resistance were induced. Antigen binding activity of the anti-CCR4 chimeric antibody in culture supernatants in wells in which growth of transformants was observed was measured by the ELISA described in the item 2 of Example 8.

Regarding the transformants in wells in which production of the anti-CCR4 chimeric antibody was observed in culture supernatants, the MTX concentration was increased by the same method, and a transformant capable of growing in the Hybridoma-SFM-FBS(5) medium comprising 200 nM MTX and of producing the anti-CCR4 chimeric antibody in a large amount was finally obtained. The obtained transformant was made into a single cell (cloning) by limiting dilution twice, and the obtained cloned cell line was named KM2760 #58-35-16. In this case, using the method for determining the transcription product of an α -1,6-fucosyltransferase gene shown in Example 9, a cell line

producing a relatively small amount of the transcription product was selected and used as a suitable cell line.

(2) Preparation of antibody-producing cell using CHO/DG44 cell

After introducing 4 μ g of the anti-CCR4 chimeric antibody expression vector pKANTEX2160 into 1.6×10^6 cells of CHO/DG44 cell by electroporation [Cytotechnology, 3, 133 (1990)], the cells were suspended in 10 ml of IMDM-dFBS(10)-HT(1) [IMDM medium (manufactured by Invitrogen) comprising 10% dFBS (manufactured by Invitrogen) and 1 x concentration of HT supplement (manufactured by Invitrogen)] and dispensed in 100 μ l/well into a 96 well culture plate (manufactured by Iwaki Glass). After culturing them at 37°C for 24 hours in a 5% CO₂ incubator, the medium was changed to IMDM-dFBS(10) (IMDM medium comprising 10% of dialyzed FBS), followed by culturing for 1 to 2 weeks. Culture supernatant was recovered from wells in which the growth was observed due to formation of a transformant showing HT-independent growth, and an expression level of the anti-CCR4 chimeric antibody in the supernatant was measured by the ELISA described in the item 2 of Example 8.

Regarding the transformants in wells in which production of the anti-CCR4 chimeric antibody was observed in culture supernatants, in order to increase an amount of the antibody production using a DHFR gene amplification system, each of them was suspended in the IMDM-dFBS(10) medium comprising 50 nM MTX to give a density of 1 to 2×10^5 cells/ml, and the suspension was dispensed in 0.5 ml into wells of a 24 well plate (manufactured by Iwaki Glass). After culturing them at 37°C for 1 to 2 weeks in a 5% CO₂ incubator, transformants showing 50 nM MTX resistance were induced. Regarding the transformants in wells in which the growth was observed, the MTX

concentration was increased to 200 nM by the same method, and a transformant capable of growing in the IMDM-dFBS(10) medium comprising 200 nM MTX and of producing the anti-CCR4 chimeric antibody in a large amount was finally obtained. The obtained transformant was named 5-03. In this case, using the method for determining the transcription product of an α -1,6-fucosyltransferase gene shown in Example 9, a cell line producing a relatively small amount of the transcription product was selected and used as a suitable cell line.

2. Antibody binding activity to CCR4 partial peptide (ELISA)

Compound 1 (SEQ ID NO:25) was selected as a human CCR4 extracellular region peptide capable of reacting with the anti-CCR4 chimeric antibody. In order to use it in the activity measurement by ELISA, a conjugate with BSA (bovine serum albumin) (manufactured by Nakalai Tesque) was prepared by the following method and used as the antigen. That is, 100 ml of a DMSO solution comprising 25 mg/ml SMCC [4-(*N*-maleimidomethyl)-cyclohexane-1-carboxylic acid *N*-hydroxysuccinimide ester] (manufactured by Sigma) was added dropwise to 900 ml of a 10 mg BSA-containing PBS solution under stirring using a vortex, followed by gently stirring for 30 minutes. A 1 ml portion of the reaction solution was applied to a gel filtration column such as NAP-10 column or the like equilibrated with 25 ml of PBS, and then eluted with 1.5 ml of PBS and the resulting eluate was used as a BSA-SMCC solution (BSA concentration was calculated based on A_{280} measurement). Next, 250 ml of PBS was added to 0.5 mg of Compound 1 and then completely dissolved by adding 250 ml of DMF, and the BSA-SMCC solution was added thereto under vortex, followed by gently stirring for 3 hours. The reaction solution was dialyzed against PBS at 4°C overnight, sodium azide was added

thereto to give a final concentration of 0.05%, and the mixture was filtered through a 0.22 mm filter to be used as a BSA-compound 1 solution.

The prepared conjugate was dispensed at 0.05 µg/ml and 50 µl/well into a 96 well EIA plate (manufactured by Greiner) and incubated for adhesion at 4°C overnight. After washing each well with PBS, 1% BSA-PBS was added thereto in 100 µl/well and allowed to react at room temperature to block the remaining active groups. After washing each well with PBS containing 0.05% Tween 20 (hereinafter referred to as "Tween-PBS"), a culture supernatant of a transformant was added at 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction, each well was washed with Tween-PBS, and then a peroxidase-labeled goat anti-human IgG(γ) antibody solution (manufactured by American Qualex) diluted 6000 times with 1% BSA-PBS as the secondary antibody was added at 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, the ABTS substrate solution was added at 50 µl/well for color development, and 20 minutes thereafter, the reaction was stopped by adding a 5% SDS solution at 50 µl/well. Thereafter, the absorbance at OD₄₁₅ was measured. The anti-CCR4 chimeric antibody obtained in the item 1 of Example 8 showed the binding activity to CCR4.

3. Purification of anti-CCR4 chimeric antibody

(1) Culturing of antibody-producing cell derived from YB2/0 cell and purification of antibody

The anti-CCR4 chimeric antibody-expressing transformant cell clone KM2760 # 58-35-16 obtained in the item 1(1) of Example 8 was suspended in Hybridoma-SFM (manufactured by Invitrogen) medium comprising 200 nM MTX and 5% of Daigo's GF21 (manufactured by Wako Pure Chemical Industries) to give a density of 2×10^5 cells/ml and

subjected to fed-batch shaking culturing using a spinner bottle (manufactured by Iwaki Glass) in a constant temperature chamber of 37°C. After culturing them for 8 to 10 days, the anti-CCR4 chimeric antibody was purified from the culture supernatant recovered using Prosep-A (manufactured by Millipore) column and gel filtration. The purified anti-CCR4 chimeric antibody was named KM2760-1.

(2) Culturing of antibody-producing cell derived from CHO-DG44 cell and purification of antibody

The anti-CCR4 chimeric antibody-producing transformant cell line 5-03 obtained in the item 1(2) of Example 8 was cultured at 37°C in a 5% CO₂ incubator using IMDM-dFBS(10) medium in a 182 cm² flask (manufactured by Greiner). When the cell density reached confluent after several days, the culture supernatant was discarded, and the cells were washed with 25 ml of PBS buffer and then mixed with 35 ml of EXCELL 301 medium (manufactured by JRH). After culturing them at 37°C for 7 days in a 5% CO₂ incubator, the culture supernatant was recovered. The anti-CCR4 chimeric antibody was purified from the culture supernatant using Prosep-A (manufactured by Millipore) column in accordance with the manufacture's instructions. The purified anti-CCR4 chimeric antibody was named KM3060.

When the binding activity to CCR4 of KM2760-1 and KM3060 was measured by ELISA, they showed equivalent binding activity.

4. Analysis of purified anti-CCR4 chimeric antibodies

Each (4 µg) of the two kinds of the anti-CCR4 chimeric antibodies produced by and purified from in different animal cells, obtained in the item 1 of this Example was subjected to SDS-PAGE in accordance with a known method [*Nature*, 227, 680 (1970)], and the molecular weight and purification degree were analyzed. In each of

the purified anti-CCR4 chimeric antibodies, a single band corresponding to the molecular weight of about 150 Kd was found under non-reducing conditions, and two bands of about 50 Kd and about 25 Kd were found under reducing conditions. The molecular weights almost coincided with the molecular weights deduced from the cDNA nucleotide sequences of antibody H chain and L chain (H chain: about 49 Kd, L chain: about 23 Kd, whole molecule: about 144 Kd) and coincided with reports stating that an IgG type antibody has a molecular weight of about 150 Kd under non-reducing conditions and is degraded into H chain having a molecular weight of about 50 Kd and L chain having a molecular weight of about 25 Kd under reducing conditions caused by cutting an S-S bond in the molecule [Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Chapter 14 (1988), Monoclonal Antibodies: Principles and Practice, Academic Press Limited (1996)], thus confirming that the anti-CCR4 chimeric antibody was expressed and purified as an antibody molecule having a correct structure.

5. Preparation of anti-CCR4 chimeric antibody having different ratio of α -1,6-fucose-free sugar chain

Sugar chains which are bound to anti-CCR4 chimeric antibody KM2760-1 derived from YB2/0 cell prepared in the item 3(1) of Example 8 and the anti-CCR4 chimeric antibody KM3060 derived from CHO/DG44 cell prepared in the item 3(2) of Example 8 were analyzed in accordance with the method of Example 10(6). The ratio of α -1,6-fucose-free sugar chains was 87% and 8% in KM2760 and KM3060, respectively. Herein, the samples are referred to as anti-CCR4 chimeric antibody (87%) and anti-CCR4 chimeric antibody (8%).

The anti-CCR4 chimeric antibody (87%) and anti-CCR4 chimeric antibody (8%) were mixed at a ratio of anti-CCR4 chimeric antibody (87%) : anti-CCR4 chimeric antibody (8%) = 1 : 39, 16 : 67, 22 : 57, 32 : 47 or 42 : 37. Sugar

Example 9

Determination of transcription product of α -1,6-fucosyltransferase gene in host cell line:

(1) Preparation of single-stranded cDNA from various cell lines

Single-stranded cDNA samples were prepared from dihydrofolate reductase gene (dhfr)-deleted CHO/DG44 cells derived from Chinese hamster ovary and rat myeloma YB2/0 cells by the following procedure.

The CHO/DG44 cells were suspended in IMDM medium (manufactured by Life Technologies) supplemented with 10% fetal bovine serum (manufactured by Life Technologies) and 1 x concentration HT supplement (manufactured by Life Technologies), and 15 ml of the suspension was inoculated into T75 flask for adhesion cell culture use (manufactured by Greiner) at a density of 2×10^5 cells/ml. Also, the YB2/0 cells were suspended in RPMI 1640 medium (manufactured by Life Technologies) supplemented with 10% fetal bovine serum (manufactured by Life Technologies) and 4 mmol/l L-GLN (manufactured by Life Technologies), and 15 ml of the suspension was inoculated into T75 flask for suspension cell culture (manufactured by Greiner) at a density of 2×10^5 cells/ml. They were cultured at 37°C in a 5% CO₂ incubator, and 1×10^7 of respective host cells were recovered on the 1st, 2nd, 3rd, 4th and 5th days of the culturing to extract total RNA using RNAeasy (manufactured by QIAGEN) in accordance with the manufacture's instructions.

The total RNA was dissolved in 45 μ l of sterile water, 1 μ l of RQ1 RNase-Free DNase (manufactured by Promega), 5 μ l of the attached 10 x DNase buffer and 0.5 μ l of RNasin Ribonuclease Inhibitor (manufactured by Promega) were added thereto, followed by reaction at 37°C for 30 minutes to degrade genome DNA contaminated in the sample. After the reaction, the total RNA was purified again using

heating at 94°C for 30 seconds, 55°C for 30 seconds and 72°C for 2 minutes as one cycle, and final heating at 72°C for 10 minutes.

After the PCR, the reaction solution was subjected to 0.8% agarose gel electrophoresis, and a specific amplified fragment of 979 bp was purified using GENECLAN Spin Kit (manufactured by BIO 101) and eluted with 10 µl of sterile water (hereinafter, the method was used for the purification of DNA fragments from agarose gel). Into a plasmid pCR2.1, 4 µl of the amplified fragment was employed to insert in accordance with the manufacture's instructions of TOPO TA Cloning Kit (manufactured by Invitrogen), and *E. coli* XL1-Blue was transformed with the reaction solution by the method of Cohen et al. [*Proc. Natl. Acad. Sci. USA*, 69, 2110 (1972)] (hereinafter, the method was used for the transformation of *E. coli*). Plasmid DNA samples were isolated in accordance with a known method [*Nucleic Acids Research*, 7, 1513 (1979)] (hereinafter, the method was used for the isolation of plasmid) from cDNA-inserted 6 clones among the obtained kanamycin-resistant colonies.

The nucleotide sequence of each cDNA inserted into the plasmid was determined using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction kit (manufactured by Parkin Elmer) in accordance with the method of the manufacture's instructions. It was confirmed that all of the inserted cDNAs of which sequences were determined by the method encode the open reading frame (ORF) partial sequences of Chinese hamster FUT8 or rat FUT8 (shown in SEQ ID NOs:6 and 7). Among these, plasmid DNA samples containing absolutely no reading error by the PCR in the sequences were selected. Herein, these plasmids are referred to as CHFUT8-pCR2.1 and YBFUT8-pCR2.1.

Separately, 3 µg of a plasmid pBluescript II KS(+) (manufactured by Stratagene) was dissolved in 35 µl of NEBuffer 2 (manufactured by New England Biolabs), and 16 units of a restriction enzyme *EcoRV* (manufactured by Takara

Shuzo) were added thereto for digestion reaction at 37°C for 3 hours. To the reaction solution, 35 µl of 1 mol/l Tris-HCl buffer (pH 8.0) and 3.5 µl of *E. coli* C15-derived alkaline phosphatase (manufactured by Takara Shuzo) were added thereto, followed by reaction at 65°C for 30 minutes to thereby dephosphorylate the DNA terminus. The reaction solution was extracted with phenol/chloroform, followed by ethanol precipitation, and the recovered DNA fragment was dissolved in 100 µl of sterile water.

Each 4 µl of the amplified fragment prepared from Chinese hamster cDNA or the amplified fragment (1192 bp) prepared from rat cDNA was mixed with 1 µl of the *EcoRV*-*EcoRV* fragment (about 3.0 Kb) prepared from plasmid pBluescript II KS(+) and 5 µl of Ligation High (manufactured by Toyobo) for ligation reaction at 16°C for 30 minutes. Using the reaction solution, *E. coli* XL1-Blue was transformed, and plasmid DNA samples were isolated respectively in accordance with a known method from the obtained ampicillin-resistant clones.

The nucleotide sequence of each cDNA inserted into the plasmid was determined using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction kit (manufactured by Parkin Elmer) in accordance with the method of the manufacture's instructions. It was confirmed that all of the inserted cDNAs of which sequences were determined by the method encode the ORF full sequences of Chinese hamster β -actin or rat β -actin. Among these, plasmid DNA samples containing absolutely no reading error of bases by the PCR in the sequences were selected. Herein, the plasmids are called CHAC-pBS and YBAC-pBS.

(4) Preparation of FUT8 standard and internal control

In order to measure a transcription level of FUT8 gene mRNA in each cell, CHFT8-pCR2.1 or YBFT8-pCR2.1, as

plasmids in which cDNA partial fragments prepared in the item (2) from Chinese hamster FUT8 or rat FUT8 were inserted into pCR2.1, respectively, were digested with a restriction enzyme *EcoRI*, and the obtained linear DNAs were used as the standards for the preparation of a calibration curve. CHFT8d-pCR2.1 and YBFT8d-pCR2.1, which were obtained from the CHFT8-pCR2.1 and YBFT8-pCR2.1, by deleting 203 bp between *ScaI* and *HindIII*, an inner nucleotide sequence of Chinese hamster FUT8 and rat FUT8, respectively, were digested with a restriction enzyme *EcoRI*, and the obtained linear DNAs were used as the internal standards for FUT8 amount determination. Details thereof are described below.

Chinese hamster FUT8 and rat FUT8 standards were prepared as follows. In 40 μ l of NEBuffer 2 (manufactured by New England Biolabs), 2 μ g of the plasmid CHFT8-pCR2.1 was dissolved, 24 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. Separately, 2 μ g of the plasmid YBFT8-pCR2.1 was dissolved in 40 μ l of NEBuffer 2 (manufactured by New England Biolabs), and 24 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. By subjecting a portion of each of the reaction solutions to 0.8% agarose gel electrophoresis, it was confirmed that an *EcoRI*-*EcoRI* fragment (about 1 Kb) containing each of cDNA partial fragments of Chinese hamster FUT8 and rat FUT8 was separated from the plasmids CHFT8-pCR2.1 and YBFT8-pCR2.1 by the restriction enzyme digestion reactions. Each of the reaction solutions was diluted with 1 μ g/ml of baker's yeast t-RNA (manufactured by SIGMA) to give a concentration of 0.02 fg/ μ l, 0.2 fg/ μ l, 1 fg/ μ l, 2 fg/ μ l, 10 fg/ μ l, 20 fg/ μ l and 100 fg/ μ l and used as the Chinese hamster FUT8 and rat FUT8 standards.

EcoRI (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. Separately, 2 µg of the plasmid YBFT8d-pCR2.1 was dissolved in 40 µl of NEBuffer 2 (manufactured by New England Biolabs), and 24 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. A portion of each of the reaction solutions was subjected to 0.8% agarose gel electrophoresis, and it was confirmed that an *EcoRI*-*EcoRI* fragment (about 800 bp) containing a fragment from which 203 bp of the inner nucleotide sequences of Chinese hamster FUT8 or rat FUT8 partial fragments was deleted was separated from the plasmids CHFT8d-pCR2.1 or YBFT8d-pCR2.1 by the restriction enzyme digestion reactions. Dilutions of 2 fg/µl were prepared from the reaction solutions using 1 µg/ml baker's yeast t-RNA (manufactured by SIGMA) and used as the Chinese hamster FUT8 or rat FUT8 internal controls.

(5) Preparation of β -actin standard and internal control

In order to measure the transcription amount of β -actin gene mRNA in various host cells, CHAc-pBS and YBAC-pBS, as plasmids in which the ORF full length of each cDNA of Chinese hamster β -actin and rat β -actin prepared in the item (3) was inserted into pBluescript II KS(+), respectively, were digested with restriction enzymes *HindIII* and *PstI* and restriction enzymes *HindIII* and *KpnI*, respectively, and the digested linear DNAs were used as the standards for the preparation of a calibration curve. CHAc-d-pBS and YBAC-d-pBS which were obtained from the CHAc-pBS and YBAC-pBS by deleting 180 bp between *DraIII* and *DraIII* of an inner nucleotide sequence of Chinese hamster β -actin and rat β -actin were digested with restriction enzymes *HindIII* and *PstI* and restriction enzymes *HindIII* and *KpnI*, respectively, and the digested linear DNAs were

used as the internal standards for β -actin amount determination. Details thereof are described below.

Chinese hamster β -actin and rat β -actin standards were prepared as follows. In 40 μ l of NEBuffer 2 (manufactured by New England Biolabs), 2 μ g of the plasmid CHAC-pBS was dissolved, and 25 units of a restriction enzyme *Hind*III (manufactured by Takara Shuzo) and 20 units of *Pst*I (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. Separately, 2 μ g of the plasmid YBAC-pBS was dissolved in 40 μ l of NEBuffer 2 (manufactured by New England Biolabs), and 25 units of a restriction enzyme *Hind*III (manufactured by Takara Shuzo) and 25 units of *Kpn*I (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. A portion of each of the reaction solutions was subjected to 0.8% agarose gel electrophoresis, and it was confirmed that a *Hind*III-*Pst*I fragment and a *Hind*III-*Kpn*I fragment (about 1.2 Kb) containing the full length ORF of each cDNA of Chinese hamster β -actin and rat β -actin were separated from the plasmids CHAC-pBS and YBAC-pBS by the restriction enzyme digestion reactions. Each of the reaction solutions was diluted with 1 μ g/ml baker's yeast t-RNA (manufactured by SIGMA) to give a concentration 2 pg/ μ l, 1 pg/ μ l, 200 fg/ μ l, 100 fg/ μ l and 20 fg/ μ l and used as the Chinese hamster β -actin and or β -actin standards.

Chinese hamster β -actin and rat β -actin internal standards were prepared as follows (Fig. 23). In 100 μ l of NEBuffer 3 (manufactured by New England Biolabs) containing 100 ng/ μ l of BSA (manufactured by New England Biolabs), 2 μ g of CHAC-pBS was dissolved, and 10 units of a restriction enzyme *Dra*III (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 3 hours. DNA fragments were recovered from the reaction solution by ethanol precipitation and the DNA termini were

changed to blunt ends using DNA Blunting Kit (manufactured by Takara Shuzo) in accordance with the manufacture's instructions, and then the reaction solution was divided into two equal parts. First, to one part of the reaction solution, 35 μ l of 1 mol/l Tris-HCl buffer (pH 8.0) and 3.5 μ l of *E. coli* C15-derived alkaline phosphatase (manufactured by Takara Shuzo) were added thereto, followed by reaction at 65°C for 30 minutes for dephosphorylating the DNA termini. The DNA fragment was recovered by carrying out dephosphorylation treatment, phenol/chloroform extraction treatment and ethanol precipitation treatment and then dissolved in 10 μ l of sterile water. The remaining part of the reaction solution was subjected to 0.8% agarose gel electrophoresis to purify a DNA fragment of about 1.1 Kb containing the ORF partial fragment of Chinese hamster β -actin.

The dephosphorylated *Dra*III-*Dra*III fragment (4.5 μ l), 4.5 μ l of the *Dra*III-*Dra*III fragment of about 1.1 Kb and 5 μ l of Ligation High (manufactured by Toyobo) were mixed, followed by ligation reaction at 16°C for 30 minutes. Using the reaction solution, *E. coli* DH5 α was transformed, and plasmid DNAs were isolated in accordance with a known method from the obtained ampicillin-resistant clones. The nucleotide sequence of each plasmid DNA was determined using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction kit (manufactured by Parkin Elmer), and it was confirmed that a Chinese hamster β -actin *Dra*III-*Dra*III 180 bp inserted into the plasmid was deleted. The plasmid is referred to as CHAc-d-pBS.

Also, a plasmid in which rat β -actin *Dra*III-*Dra*III 180 bp was deleted was prepared via the same steps of CHAc-d-pBS. The plasmid is referred to as YBAc-d-pBS.

Next, 2 μ g of the plasmid CHAc-d-pBS was dissolved in 40 μ l of NEBuffer 2 (manufactured by New England

(6) Determination of transcription amount by competitive PCR

- 166 -

reaction of cDNA in each host cell line. That is, the PCR was carried out using the β -actin internal control DNA prepared in the item (5) and the host cell-derived cDNA obtained in the item (1) as the templates, the determined value of the β -actin transcription product in the host cell line was calculated from the relative value of the amount of the amplified product derived from each template. Details thereof are described below.

The FUT8 transcription product was determined by the following procedure. First, a set of sequence-specific primers (shown in SEQ ID NOS:13 and 14) common to the inner sequences of the ORF partial sequences of Chinese hamster FUT8 and rat FUT8 obtained in the item (2) were designed.

Next, PCR was carried out using a DNA polymerase ExTaq (manufactured by Takara Shuzo) in 20 μ l in total volume of a reaction solution [ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs, 0.5 μ mol/l gene-specific primers (SEQ ID NOS:13 and 14) and 5% DMSO] containing 5 μ l of 50 folds-diluted cDNA solution prepared from each of respective host cell line in the item (1) and 5 μ l (10 fg) of the plasmid for internal control. The PCR was carried out by heating at 94°C for 3 minutes and subsequent 32 cycles of heating at 94°C for 1 minute, 60°C for 1 minute and 72°C for 1 minute as one cycle.

Also, PCR was carried out in a series of reaction in which 5 μ l (0.1 fg, 1 fg, 5 fg, 10 fg, 50 fg, 100 fg, 500 fg or 1 pg) of the FUT8 standard plasmid obtained in the item (4) was added instead of the each host cell line-derived cDNA, and used in the preparation of a calibration curve for the FUT8 transcription level.

The β -actin transcription product was determined by the following procedure. First, two sets of respective gene-specific primers common to the inner sequences of the ORF full lengths of Chinese hamster β -actin and rat β -actin obtained in the item (3) were designed (the former are

shown in SEQ ID NOs:15 and 16, and the latter are shown in SEQ ID NOs:17 and 18).

Next, PCR was carried out using a DNA polymerase ExTaq (manufactured by Takara Shuzo) in 20 μ l in total volume of a reaction solution [ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs, 0.5 μ mol/l gene-specific primers (SEQ ID NOs:15 and 16, or SEQ ID NOs:17 and 18) and 5% DMSO] containing 5 μ l of 50 folds-diluted cDNA solution prepared from respective host cell line in the item (1) and 5 μ l (1 pg) of the plasmid for internal control. The PCR was carried out by heating at 94°C for 3 minutes and subsequent 17 cycles of heating at 94°C for 30 seconds, 65°C for 1 minute and 72°C for 2 minutes as one cycle.

Also, PCR was carried out in a series of reaction in which 5 μ l (10 pg, 5 pg, 1 pg, 500 fg or 100 fg) of the β -actin standard plasmid obtained in the item (5) was added instead of the each host cell line-derived cDNA, and used in the preparation of a calibration curve for the β -actin transcription level.

Table 3

Target gene	Primer set *	Size (bp) of PCR amplification product	
		Target	Competitor
FUT8	F: 5'-GTCCATGGTGATCCTGCAGTGTGG-3' R: 5'-CACCAATGATATCTCCAGGTTCC-3'	638	431
β -Actin (Chinese hamster)	F: 5'-GATATCGCTGCGCTCGTTGTCGAC-3' R: 5'-CAGGAAGGAAGGCTGGAAGAGAGC-3'	789	609
β -Actin (Rat)	F: 5'-GATATCGCTGCGCTCGTCGTCGAC-3' R: 5'-CAGGAAGGAAGGCTGGAAGAGAGC-3'	789	609

* F: forward primer, R: reverse primer

By carrying out PCR using the primer set described in Table 3, a DNA fragment having a size shown in the

and 15 ml of the suspension was inoculated into T75 flask for suspension cell culture use (manufactured by Greiner) at a density of 2×10^5 cells/ml. Also, cells of the 61-33 were suspended in Hybridoma-SFM medium (manufactured by Life Technologies) supplemented with 0.2% bovine serum albumin fraction V (manufactured by Life Technologies) (hereinafter referred to as "BSA"), and 15 ml of the suspension was inoculated into T75 flask for suspension cell culture (manufactured by Greiner) at a density of 2×10^5 cells/ml. They were cultured at 37°C in a 5% CO₂ incubator, and 1, 2, 3, 4 and 5 days after culturing, 1×10^7 of respective host cells were recovered to extract total RNA using RNAeasy (manufactured by QIAGEN) in accordance with the manufacture's instructions.

The total RNA was dissolved in 45 µl of sterile water, and 1 µl of RQ1 RNase-Free DNase (manufactured by Promega), 5 µl of the attached 10 x DNase buffer and 0.5 µl of RNasin Ribonuclease Inhibitor (manufactured by Promega) were added thereto, followed by reaction at 37°C for 30 minutes to degrade genome DNA contaminated in the sample. After the reaction, the total RNA was purified again using RNAeasy (manufactured by QIAGEN) and dissolved in 50 µl of sterile water.

In a 20 µl reaction mixture using oligo(dT) as a primer, single-stranded cDNA was synthesized from 3 µg of each of the obtained total RNA samples by reverse transcription reaction using SUPERScript™ Preamplification System for First Strand cDNA Synthesis (manufactured by Life Technologies) in accordance with the manufacture's instructions. The reaction solution was diluted 50 folds with water and stored at -80°C until use.

(2) Determination of transcription amounts of each gene by competitive PCR

The transcription amount of each of the genes on the cDNA derived from the antibody-producing cell line obtained in the item (1) was determined by competitive PCR in accordance with Example 9(6).

The FUT8 gene-derived mRNA transcription amount in each of the antibody-producing cell lines was determined by the following procedure.

CHFT8-pCR2.1 and YBFT8-pCR2.1, as plasmids in which cDNA partial fragments prepared in Example 9(2) from Chinese hamster FUT8 and rat FUT8, respectively, were inserted into pCR2.1, were digested with a restriction enzyme *EcoRI*, and the obtained linear DNAs were used as the standards in the preparation of a calibration curve for determining the FUT8 transcription amount.

CHFT8d-pCR2.1 and YBFT8d-pCR2.1, which were obtained by deleting 203 bp between *ScaI* and *HindIII* of an inner nucleotide sequence of Chinese hamster FUT8 and rat FUT8, respectively, in Example 9(4) were digested with a restriction enzyme *EcoRI*, and the obtained linear DNAs were used as the internal standards for FUT8 amount determination.

PCR was carried out using a DNA polymerase ExTaq (manufactured by Takara Shuzo) in 20 μ l in total volume of a reaction solution [ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs, 0.5 μ mol/l FUT8 gene-specific primers (SEQ ID NOS:13 and 14) and 5% DMSO] containing 5 μ l of 50 folds-diluted cDNA solution prepared from each of the antibody-producing cell line in the item (1) and 5 μ l (10 fg) of the plasmid for internal control. The PCR was carried out by heating at 94°C for 3 minutes and subsequent 32 cycles of heating at 94°C for 1 minute, 60°C for 1 minute and 72°C for 1 minute as one cycle.

antibody-producing cell lines and 5 μ l (1 pg) of the plasmid for internal control. The PCR was carried out by heating at 94°C for 3 minutes and subsequent 17 cycles of heating at 94°C for 30 seconds, 65°C for 1 minute and 72°C for 2 minutes as one cycle. Also, PCR was carried out in a series of reaction in which 10 pg, 5 pg, 1 pg, 500 fg or 100 fg of the β -actin standard plasmid was added instead of the each antibody-producing cell line-derived cDNA, and used in the preparation of a calibration curve for the β -actin transcription amount. In this case, 1 μ g/ml of a baker's yeast t-RNA (manufactured by SIGMA) was used for the dilution of standard plasmid.

By PCR using the primer set described in Table 3, a DNA fragment having a size shown in the target column of Table 3 can be amplified from each gene transcription product and each standard, and a DNA fragment having a size shown in the competitor column of Table 3 can be amplified from each internal control.

A 7 μ l portion of each of the solutions after PCR was subjected to 1.75% agarose gel electrophoresis, and then the gel was stained by soaking it for 30 minutes in 1 \times concentration SYBR Green I Nucleic Acid Gel Stain (manufactured by Molecular Probes). The amount of the amplified DNA fragment was measured by calculating luminescence intensity of each amplified DNA using a fluoro-imager (FluorImager SI; manufactured by Molecular Dynamics).

The amount of the amplified product formed by PCR which used a standard plasmid as the template was measured by the method, and a calibration curve was prepared by plotting the measured values against the amounts of the standard plasmid. Using the calibration curve, the amount of cDNA of a gene of interest in each cell line was calculated from the amount of the amplified product when each antibody-producing cell line-derived cDNA was used as

the template, and the value was defined as the mRNA transcription amount in each cell line.

The FUT8 transcription amounts are shown in Table 5 as relative values to the amount of the β -actin transcription product. Throughout the culturing period, the FUT8 transcription amount in the YB2/0 cell-derived antibody-producing 61-33 was 0.3% or less of β -actin while it was 0.7% to 1.5% in the CHO cell-derived antibody-producing cell.

The results shows that the amount of the FUT8 transcription product in the YB2/0 cell-derived antibody-producing cell line was significantly less than that in the antibody-producing cell line derived from the CHO cell.

Table 5

Cell line	Culture days				
	1st	2nd	3rd	4th	5th
DCHI01-20	0.75	0.73	0.99	1.31	1.36
61-33	0.16	0.19	0.24	0.30	<0.10

Example 11

Preparation of mouse α -1,6-fucosyltransferase (FUT8) gene over-expressing cell line:

(1) Construction of mouse α -1,6-fucosyltransferase (FUT8) expression plasmid

Total RNA was extracted from 1×10^7 cells of a mouse myeloma NSO cell (RCB0213, Cell Bank at The Institute of Physical and Chemical Research) subcultured using IMDM medium (manufactured by Life Technologies) containing 10% fetal bovine serum (manufactured by Life Technologies), using RNAeasy (manufactured by QIAGEN) in accordance with the manufacture's instructions. The total RNA was dissolved in 45 μ l of sterile water, and 1 μ l of RQ1 RNase-Free DNase (manufactured by Promega), 5 μ l of the attached 10 \times DNase buffer and 0.5 μ l of RNasin Ribonuclease

E. coli DH5 α was transformed with the reaction solution. Plasmid DNAs were isolated in accordance with a known method from cDNA-inserted 6 clones among the obtained kanamycin-resistant colonies.

The nucleotide sequence of each cDNA inserted into the plasmid was determined using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction kit (manufactured by Parkin Elmer) in accordance with the method of the manufacture's instructions. It was confirmed that all of the inserted cDNAs of which sequences were determined encode the ORF full sequence of mouse FUT8. Among these, a plasmid DNA containing absolutely no reading error of bases by the PCR in the sequences were selected (its DNA sequence and amino acid sequence are shown in SEQ ID NOS:2 and 24, respectively). Also, inconsistency of 3 bases due to amino acid substitution was found in the sequence when compared with the mouse FUT8 sequence registered on GenBank. Herein, the plasmid is referred to mFUT8-pCR2.1.

Next, a plasmid pBSmFUT8 containing the ORF full sequence of mouse FUT8 was constructed as follows (Fig. 26). First, 1 μ g of a plasmid pBluescript II KS(+) (manufactured by Stratagene) was dissolved in 35 μ l of NEBuffer 2 (manufactured by New England Biolabs), and 20 units of a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 2 hours. To the reaction solution, 35 μ l of 1 mol/l Tris-HCl buffer (pH 8.0) and 3.5 μ l of *E. coli* C15-derived alkaline phosphatase (manufactured by Takara Shuzo) were added, followed by reaction at 65°C for 30 minutes for dephosphorylate the DNA termini. The reaction solution was extracted with phenol/chloroform, followed by ethanol precipitated, and the recovered DNA fragment was dissolved in 10 μ l of sterile water.

μ l of 1 mol/l Tris-HCl buffer (pH 8.0) and 3.5 μ l of *E. coli* C15-derived alkaline phosphatase (manufactured by Takara Shuzo) were added thereto, followed by reaction at 65°C for 30 minutes to dephosphorylate the DNA termini. The reaction solution was extracted with phenol/chloroform extraction, followed by ethanol precipitation, and the recovered DNA fragment was dissolved in 10 μ l of sterile water.

Separately, 1 μ g of pBSmfFUT8 was dissolved in 50 μ l of Universal Buffer H (manufactured by Takara Shuzo), and 20 units of a restriction enzyme *Sal*I (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. A DNA fragment was recovered from the reaction solution by ethanol precipitation and dissolved in 35 μ l of NEBuffer 2 (manufactured by New England Biolabs), and 20 units of a restriction enzyme *Bam*HI (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, the solution was subjected to 0.8% agarose gel electrophoresis to purify a DNA fragment of about 1.7 Kb containing the ORF full sequence of mouse FUT8 cDNA.

The obtained plasmid pAGE249-derived *Bam*HI-*Sal*I fragment (1 μ l, 6.5 Kb), 4 μ l of the *Bam*HI-*Sal*I fragment (1.7 Kb) prepared from the plasmid pBSmfFUT8 and 5 μ l of Ligation High (manufactured by Toyobo) were mixed, followed by ligation reaction at 16°C for 30 minutes. Using the reaction solution, *E. coli* DH5 α was transformed, and a plasmid DNA was isolated in accordance with a known method from the obtained ampicillin-resistant clones. Herein, the plasmid is referred to pAGEmfFUT8.

(2) Preparation of mouse α -1,6-fucosyltransferase (FUT8) gene over-expressing cell line

A stable FUT8 gene-expressing cell line was obtained by introducing the mouse FUT8 expression vector pAGEMfFUT8 constructed in the item (1) into 61-33. The 61-33 is a clone obtained by carrying out serum-free adaptation of a transformant cell 7-9-51 (FERM BP-6691, International Patent Organism Depositary, National Institute of Advanced Industrial Science and Technology) derived from a YB2/0 cell highly producing an anti-ganglioside GD3 chimeric antibody, and then carrying out single cell isolation by two limiting dilution.

The plasmid pAGEMfFUT8 was transferred into 61-33 by the following procedure in accordance with the electroporation [Cytotechnology, 3, 133 (1990)]. First, 30 μ g of the plasmid pAGEMfFUT8 was dissolved in 600 μ l of NEBuffer 4 (manufactured by New England Biolabs), and 100 units of a restriction enzyme *Fsp*I (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours to obtain a linear fragment. The reaction solution was subjected to ethanol precipitation, and the recovered linear plasmid was made into a 1 μ g/ μ l aqueous solution. Next, the 61-33 was suspended in a K-PBS buffer (137 mol/l KCl, 2.7 mol/l NaCl, 8.1 mol/l Na_2HPO_4 , 1.5 mol/l KH_2PO_4 , 4.0 mol/l MgCl_2) to give a density of 2×10^7 cells/ml, and 200 μ l of the cell suspension (4×10^6 cells) was mixed with 10 μ l (10 μ g) of the linear plasmid. The cell-DNA mixture was transferred into Gene Pulser Cuvette (inter-electrode distance, 2 mm) (manufactured by BIO-RAD) and then electroporation was carried out using a cell fusion apparatus Gene Pulser (manufactured by BIO-RAD) at 0.2 KV pulse voltage and 250 μ F electric capacity. The cell suspension was mixed with 10 ml of Hybridoma-SFM medium (manufactured by Life Technologies) supplemented with 5% fetal bovine dialyzed

serum (manufactured by Life Technologies) and 0.2% BSA (manufactured by Life Technologies) and dispensed in 100 μ l portions into a 96 well plate for suspension cell use (manufactured by Greiner). After culturing them at 37°C for 24 hours in 5% CO₂, 50 μ l of the culture supernatant was removed, and Hybridoma-SFM medium (manufactured by Life Technologies) supplemented with 0.5 mg/ml Hygromycin B (manufactured by Wako Pure Chemical Industries), 5% fetal bovine dialyzed serum (manufactured by Life Technologies) and 0.2% BSA (manufactured by Life Technologies) was dispensed at 100 μ l. They were cultured for 3 weeks while repeating the medium exchange step at intervals of 3 to 4 days, and 14 cell lines showing hygromycin resistance were obtained.

On the other hand, a negative control cell line was prepared by introducing the plasmid pAGE249 as a parent vector of the pAGEMfUT8 into the 61-33. According to the above procedure, 10 μ g of the plasmid pAGE249 converted into linear form with a restriction enzyme FspI was introduced into 4×10^6 cells of the 61-33 using the electroporation. The cells were mixed with 15 ml of Hybridoma-SFM medium (manufactured by Life Technologies) supplemented with 5% fetal bovine dialyzed serum (manufactured by Life Technologies) and 0.2% BSA (manufactured by Life Technologies), transferred into a T75 flask for suspension cell (manufactured by Greiner) and then cultured at 37°C for 24 hours in 5% CO₂. After culturing them, a half of the culture supernatant (7.5 ml) was removed by centrifugation at 800 rpm for 4 minutes, and the cells were suspended in 7.5 ml of Hybridoma-SFM medium (manufactured by Life Technologies) supplemented with 0.5 mg/ml Hygromycin B (manufactured by Wako Pure Chemical Industries), 5% fetal bovine dialyzed serum (manufactured by Life Technologies) and 0.2% BSA (manufactured by Life Technologies) and transferred into the T75 flask for

- 182 -

In a 20 μ l reaction mixture using oligo(dT) as a primer, single-stranded cDNA was synthesized from 2.5 μ g of the obtained total RNA by reverse transcription reaction using SUPERSCRIPTM Preamplification System for First Strand cDNA Synthesis (manufactured by Life Technologies) in accordance with the manufacture's instructions. The reaction solution was diluted 50 folds with water and the transcription amount of each gene was determined by the competitive PCR in accordance with Example 9(6).

The FUT8 gene-derived mRNA transcription amount in each expression cell line was determined by the following procedure.

YBFT8-pCR2.1, as a plasmid in which a cDNA partial fragment prepared in Example 9(2) from rat FUT8 was inserted into pCR2.1, was digested with a restriction enzyme *Eco*RI, and the obtained linear DNA was used as the standard in the preparation of a calibration curve for determining the FUT8 transcription amount.

Among the YBFT8-pCR2.1 prepared in Example 9(4), YBFT8d-pCR2.1 obtained by deleting 203 bp between *Sca*I and *Hind*III of an inner nucleotide sequence of rat FUT8 was digested with a restriction enzyme *Eco*RI, and the obtained linear DNA was used as the internal control for FUT8 determination.

PCR was carried out using a DNA polymerase ExTaq (manufactured by Takara Shuzo) in 20 μ l in total volume of a reaction solution [ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs, 0.5 μ mol/l FUT8 gene-specific primers (SEQ ID NOs:13 and 14) and 5% DMSO] containing 5 μ l of 50 folds-diluted cDNA solution prepared from respective expression cell line in the above and 5 μ l (10 fg) of the plasmid for internal control. The PCR was carried out by heating at 94°C for 3 minutes and subsequent 32 cycles of heating at 94°C for 1 minute, 60°C for 1 minute and 72°C for 1 minute as one cycle.

Also, PCR was carried out in a series of reaction in which 5 μ l (0.1 fg, 1 fg, 5 fg, 10 fg, 50 fg, 100 fg, 500 fg or 1 pg) of the FUT8 standard plasmid was added instead of the each expression cell line-derived cDNA, and used in the preparation of a calibration curve for the FUT8 transcription amount. In this case, 1 μ g/ml baker's yeast t-RNA (manufactured by SIGMA) was used for the dilution of standard plasmid.

On the other hand, since it is considered that the β -actin gene is transcribed constantly in each cell and its transcription level is approximately the same between cells, the transcription amount of the β -actin gene was determined as an index of the efficiency of synthesis reaction of cDNA in each expression cell line.

YBAC-pBS, as a plasmid in which the ORF full sequence of cDNA of rat β -actin was inserted into pBluescript II KS(+) prepared in Example 9(3), was digested with restriction enzymes *Hind*III and *Kpn*I, and the obtained linear DNA was used as the standard in the preparation of a calibration curve for determining the β -actin gene transcription amount.

YBACd-pBS obtained from the YBAC-pBS by deleting 180 bp between *DraI* and *DraI* of an inner nucleotide sequence of rat β -actin was digested with restriction enzymes *HindIII* and *KpnI*, and the obtained linear DNA was used as the internal standards for β -actin amount determination.

PCR was carried out using a DNA polymerase ExTaq (manufactured by Takara Shuzo) in 20 μ l in total volume of a reaction solution [ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs, 0.5 μ mol/l β -actin-specific primers (SEQ ID NOs:17 and 18) and 5% DMSO] containing 5 μ l of 50 folds-diluted cDNA solution prepared from each of the expression cell lines and 5 μ l (1 pg) of the plasmid for internal control. The PCR was carried out by heating at

Fig. 28 shows the FUT8 transcription amounts as relative values to the amount of β -actin transcription product. Three cell lines mfFUT8-1, mfFUT8-2 and mfFUT8-4 and the PAGE249-introduced cell line were cell lines having a relatively small FUT8 transcription amount, which was equivalent to 0.3 to 10% of a β -actin transcription amount. On the other hand, other three cell lines mfFUT8-3, mfFUT8-6 and mfFUT8-7 were cell lines having a relatively large FUT8 transcription amount, which was equivalent to 20 to 40% of a β -actin transcription amount.

(4) Purification of antibody produced by α -1,6-fucosyltransferase (FUT8) gene over-expressing cell line

Each of the six FUT8 gene over-expressing cell lines and one negative control cell line obtained in the item (2) was suspended in Hybridoma-SFM medium (manufactured by Life Technologies) supplemented with 200 nmol/l MTX, 0.5 mg/ml Hygromycin B (manufactured by Wako Pure Chemical Industries) and 0.2% BSA (manufactured by Life Technologies) to give a density of 2×10^5 cells/ml, and then 100 ml in total of the suspension was inoculated into three T225 flasks for suspension cell culture use (manufactured by IWAKI). After culturing them at 37°C for 7 to 9 days in a 5% CO₂ incubator, the number of intact cells was counted to confirm that their viability was almost the same (each 30% or less), and then each cell suspension was recovered. Each of the cell suspensions was centrifuged at 3,000 rpm at 4°C for 10 minutes, and the recovered supernatant was centrifuged at 10,000 rpm at 4°C for 1 hour and then filtered using PES Filter Unit (manufactured by NALGENE) having a pore diameter of 0.22 μ m with 150 ml capacity.

Prosep-A HighCapacity (manufactured by bioPROCESSING) was packed in a 0.8 cm diameter column to a

thickness of 2 cm and washed with 10 ml of 0.1 mol/l citrate buffer (pH 3.0) and 10 ml of 1 mol/l glycine/NaOH-0.15 mol/l NaCl buffer (pH 8.6) in that order to effect equilibration the carrier. Next, 100 ml of each of the culture supernatant was passed through the column and washed with 50 ml of 1 mol/l glycine/NaOH-0.15 mol/l NaCl buffer (pH 8.6). After washing them, the antibody absorbed to Prosep-A was eluted using 2.5 ml of a 0.1 mol/l citrate buffer (pH 3.0), the eluate was fractionated at 500 μ l and each fraction was neutralized by mixing with 100 μ l of 2 mol/l Tris-HCl (pH 8.5). Two fractions containing the antibody at a high concentration (1.2 ml in total) were selected by the BCA method [Anal. Biochem., 150, 76 (1985)], combined and then dialyzed against 10 mol/l citrate buffer (pH 6.0) at 4°C for a whole day and night. After the dialysis, the antibody solution was recovered and subjected to sterile filtration using a 0.22 μ m pore size Millex GV (manufactured by MILLIPORE).

(5) *In vitro* cytotoxic activity (ADCC activity) of antibody produced by mouse α -1,6-fucosyltransferase (FUT8) gene over-expressing cell line

In order to evaluate *in vitro* cytotoxic activity of the anti-GD3 antibodies purified in the item (4), ADCC activity was measured using a GD3-positive cell, human melanoma cultured cell line G-361 (RCB0991, Cell Bank at The Institute of Physical and Chemical Research).

The G-361 cells subcultured in RPMI1640 medium (manufactured by Life Technologies) containing 10% fetal bovine serum (manufactured by Life Technologies) (hereinafter referred to as "RPMI1640-FBS(10)") were suspended in 500 μ l of RPMI1640-FBS(10) at a density of 1×10^6 cells, and 3.7 MBq of $\text{Na}_2^{51}\text{CrO}_4$ was added thereto, followed by culturing at 37°C for 30 minutes for labeling the cells with a radioisotope. After centrifugation at

1,200 rpm for 5 minutes, the supernatant was discarded and the target cells were suspended in 5 ml of RPMI1640-FBS(10). The washing step was repeated three times and then the cell suspension was incubated for 30 minutes on ice for spontaneous dissociation of the radioactive substance. The washing step was again repeated twice and then the cells were suspended in 5 ml of RPMI1640-FBS(10) to thereby prepare 2×10^5 cells/ml of a target cell suspension.

On the other hand, 30 ml of peripheral blood was collected from a healthy person and gently mixed with 0.5 ml of heparin sodium (manufactured by Shimizu Pharmaceutical) and then mixed with 30 ml of physiological saline (manufactured by Otsuka Pharmaceutical). After the mixing, 10 ml of the mixture was gently overlaid on 4 ml of Lymphoprep (manufactured by NYCOMED PHARMA AS) and centrifuged at room temperature at 2,000 rpm for 30 minutes. The separated mononuclear cell fractions were collected from the centrifugation tubes, combined and then suspended in 30 ml of RPMI1640-FBS(10). After centrifugation at room temperature at 1,200 rpm for 5 minutes, the supernatant was discarded and the cells were suspended in 20 ml of RPMI1640-FBS(10). The washing step was repeated twice and then 2×10^6 cells/ml of an effector cell suspension was prepared using RPMI1640-FBS(10).

The target cell suspension was dispensed at 50 μ l (1×10^4 cells/well) into each well of a 96 well U-bottom plate (manufactured by Falcon). Subsequently, the effector cell suspension was dispensed at 100 μ l (2×10^5 cells/well) into each well to thereby adjust the ratio of the effector cells to the target cells to 20 : 1. Next, using a 10 M citrate buffer (pH 6.0), a series of dilution solution of 0.01 μ g/ml, 0.1 μ g/ml, 1 μ g/ml and 10 μ g/ml of each anti-GD3 antibody obtained in the item (4) was prepared, and the diluted solutions were dispensed at 50 μ l into the wells to give final concentrations of 0.0025 μ g/ml,

0.025 µg/ml, 0.25 µg/ml and 2.5 µg/ml, respectively. After carrying out the reaction at 37°C for 4 hours in 5% CO₂, the plate was centrifuged at 1,200 rpm for 5 minutes. Into a 12 mm diameter RIA tube (manufactured by IWAKI), 50 µl of the supernatant in each well was transferred and, and the amount of the dissociated ⁵¹Cr was measured using MINAX-γ auto-gamma counter 5550 (manufactured by PACKARD).


Also, the amount of the spontaneously dissociated ⁵¹Cr was calculated by carrying out the same reaction in a reaction mixture in which 150 µl of RPMI1640-FBS(10) was added instead of the effector cell suspension and antibody solution. The amount of the total dissociated ⁵¹Cr was calculated by carrying out the same reaction in a reaction mixture in which 100 µl of 1 N hydrochloric acid and 50 µl of RPMI1640-FBS(10) were added instead of the effector cell suspension and antibody solution. Using these values, the ADCC activity was calculated based on the formula (II) described in the item 2(3) of Example 2.


Fig. 29 shows ADCC activity of each of the anti-GD3 antibodies for G-361 cell. Three cell lines mfFUT8-1, mfFUT8-2 and mfFUT8-4 having a low FUT8 expression level as shown in Fig. 28 showed potent ADCC activity equivalent to that of the negative control PAGE249-introduced cell line. On the other hand, other three cell lines mfFUT8-3, mfFUT8-6 and mfFUT8-7 having a high FUT8 expression level as shown in Fig. 28 showed low ADCC activity equivalent to that of the anti-GD3 antibody produced from CHO cell. Based on these results, it was shown that the ADCC activity of produced antibodies can be controlled by regulating the expression level of FUT8 in host cells.

(6) Sugar chain analysis of antibody produced by mouse α -1,6-fucosyltransferase (FUT8) gene over-expressing cell line

Sugar chains of the anti-GD3 antibodies purified in the item (4) were analyzed. The sugar chains binding to the antibodies produced by mfFUT8-6 and pAGE249-introduced cell lines were cleaved from proteins by subjecting the antibodies to hydrazinolysis [*Method of Enzymology*, 83, 263 (1982)]. After removing hydrazine by evaporation under a reduced pressure, *N*-acetylation was carried out by adding an aqueous ammonium acetate solution and acetic anhydride. After freeze-drying, fluorescence labeling by 2-aminopyridine [*J. Biochem.*, 95, 197 (1984)] was carrying out. A fluorescence-labeled sugar chain group (PA-treated sugar chain group) was separated from excess reagents using Superdex Peptide HR 10/30 column (manufactured by Pharmacia). The sugar chain fractions were dried using a centrifugation concentrator and used as a purified PA-treated sugar chain group. Next, the purified PA-treated sugar chain group was subjected to reverse phase HPLC analysis using a CLC-ODS column (manufactured by Shimadzu) (Fig. 30). When calculated from the peak area, the content of α -1,6-fucose-free sugar chains in mfFUT8-6 was 10%, and the content of α -1,6-fucose-bound sugar chains was 90%. The content of α -1,6-fucose-free sugar chains in pAGE249 was 20%, and the content of α -1,6-fucose-bound sugar chains was 80%. Based on these results, it was found that the content of α -1,6-fucose-bound sugar chains of a produced antibody is increased by over-expressing the FUT8 gene.

Fig. 30 shows elution patterns obtained by carrying out reverse phase HPLC analysis of each of PA-treated sugar chains prepared from antibodies produced by mfFUT8-6 and pAGE249-introduced cell lines. Figs. 30A and 30B show elution patterns of mfFUT8-6 and pAGE249, respectively. The relative fluorescence intensity and the elution time





amplified fragment of about 2 Kb was purified. Into a plasmid pCR2.1, 4 μ l of the DNA fragment was employed to insert in accordance with the instructions attached to TOPO TA Cloning Kit (manufactured by Invitrogen), and *E. coli* DH5 α was transformed with the reaction solution. Plasmid DNAs were isolated in accordance with a known method from cDNA-inserted 8 clones among the obtained kanamycin-resistant colonies.

The nucleotide sequence of each cDNA inserted into the plasmid was determined using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction Kit (manufactured by Parkin Elmer) in accordance with the method of the manufacture's instructions. It was confirmed by the method that all of the inserted cDNAs encode a sequence containing the full ORF of CHO cell FUT8. Among these, a plasmid DNA containing absolutely no reading error of bases by the PCR in the sequences was selected. Herein, the plasmid is referred to as CHfFUT8-pCR2.1. The determined nucleotide sequence and the amino acid sequence of the cDNA of CHO FUT8 are shown in SEQ ID NOS:1 and 23, respectively.

(2) Preparation of CHO cell α -1,6-fucosyltransferase (FUT8) genomic sequence

Using the ORF full length cDNA fragment of CHO cell FUT8 obtained in the item (1) as a probe, a CHO cell FUT8 genomic clone was obtained in accordance with a known genome screening method described, e.g., in *Molecular Cloning, Second Edition, Current Protocols in Molecular Biology, A Laboratory Manual, Second Edition* (1989). Next, after digesting the obtained genomic clone using various restriction enzymes, the Southern hybridization was carried out using an AfaI-Sau3AI fragment (about 280 bp) containing initiation codon of the CHO cell FUT8 cDNA as a probe, and then a XbaI-XbaI fragment (about 2.5 Kb) and a SacI-SacI

In 35 μ l of NEBuffer 4 (manufactured by New England Biolabs), 1.0 μ g of a plasmid pKSelectPuro (manufactured by Lexicon) was dissolved, and 20 units of a restriction enzyme AscI (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, the solution was subjected to 0.8% (w/v) agarose gel electrophoresis to purify a DNA fragment of about 1.5 Kb containing a puromycin resistance gene expression unit.

On the other hand, 1.0 μ g of a plasmid ploxP described in Japanese Published Examined Patent Application No. 314512/99 was dissolved in 35 μ l of NEBuffer 4 (manufactured by New England Biolabs), and 20 units of a restriction enzyme AscI (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, the solution was subjected to 0.8% (w/v) agarose gel electrophoresis to purify a DNA fragment of about 2.0 Kb.

The obtained AscI-AscI fragment (4.5 μ l, about 1.5 Kb) derived from the plasmid pKSelectPuro, 0.5 μ l of the AscI-AscI fragment (about 2.0 Kb) derived from the plasmid ploxP and 5.0 μ l of Ligation High (manufactured by Toyobo) were mixed, followed by ligation reaction at 16°C for 30 minutes. *E. coli* DH5 α was transformed using the reaction solution, and a plasmid DNA was isolated in accordance with a known method from the obtained ampicillin-resistant clones. Herein, the plasmid is referred to as ploxPPuro.

(2) Construction of plasmid pKOFUT8gE2-1

A plasmid pKOFUT8gE2-1 was constructed by the following procedure (Fig. 34), using the plasmid pFUT8fgE2-2 obtained in Example 12(2) having a genome region comprising exon 2 of Chinese hamster FUT8.

In 35 μ l of NEBuffer 1 (manufactured by New England Biolabs) containing 100 μ g/ml of BSA (manufactured by New

England Biolabs), 2.0 µg of the plasmid pFUT8fgE2-2 was dissolved, and 20 units of a restriction enzyme *SacI* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. A DNA fragment was recovered from the reaction solution by ethanol precipitation and dissolved in 35 µl of NEBuffer 2 (manufactured by New England Biolabs) containing 100 µg/ml of BSA (manufactured by New England Biolabs), and 20 units of a restriction enzyme *EcoRV* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, the solution was subjected to 0.8% (w/v) agarose gel electrophoresis to purify a DNA fragment of about 1.5 Kb.

Separately, 1.0 µg of a plasmid LITMUS28 (manufactured by New England Biolabs) was dissolved in 35 µl of NEBuffer 1 (manufactured by New England Biolabs) containing 100 µg/ml of BSA (manufactured by New England Biolabs), and 20 units of a restriction enzyme *SacI* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. A DNA fragment was recovered from the reaction solution by ethanol precipitation and dissolved in 35 µl of NEBuffer 2 (manufactured by New England Biolabs) containing 100 µg/ml of BSA (manufactured by New England Biolabs), and 20 units of a restriction enzyme *EcoRV* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, the solution was subjected to 0.8% (w/v) agarose gel electrophoresis to purify a DNA fragment of about 2.8 Kb.

The obtained *EcoRV*-*SacI* fragment (4.5 µl, about 1.5 Kb) derived from the plasmid pFUT8fgE2-2, 0.5 µl of the *EcoRV*-*SacI* fragment (about 2.8 Kb) derived from the plasmid LITMUS28 and 5.0 µl of Ligation High (manufactured by Toyobo) were mixed, followed by ligation reaction at 16°C for 30 minutes. *E. coli* DH5α was transformed using the

reaction solution, and a plasmid DNA was isolated in accordance with a known method from the obtained ampicillin-resistant clones. Herein, the plasmid is referred to as pKOFUT8gE2-1.

(3) Construction of plasmid pKOFUT8gE2-2

A plasmid pKOFUT8gE2-2 was constructed by the following procedure (Fig. 35), using the plasmid pKOFUT8gE2-1 obtained in the item (2).

In 30 μ l of NEBuffer 2 (manufactured by New England Biolabs) containing 100 μ g/ml of BSA (manufactured by New England Biolabs), 2.0 μ g of the plasmid pKOFUT8gE2-1 was dissolved, and 20 units of a restriction enzyme *EcoRV* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. A DNA fragment was recovered from the reaction solution by ethanol precipitation and dissolved in 30 μ l of NEBuffer 1 (manufactured by New England Biolabs) containing 100 μ g/ml of BSA (manufactured by New England Biolabs), and 20 units of a restriction enzyme *KpnI* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, the solution was subjected to 0.8% (w/v) agarose gel electrophoresis to purify a DNA fragment of about 1.5 Kb.

Separately, 1.0 μ g of the plasmid ploxPPuro was dissolved in 30 μ l of NEBuffer 4 (manufactured by New England Biolabs), and 20 units of a restriction enzyme *HpaI* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. A DNA fragment was recovered from the reaction solution by ethanol precipitation and dissolved in 30 μ l of NEBuffer 1 (manufactured by New England Biolabs) containing 100 μ g/ml of BSA (manufactured by New England Biolabs), and 20 units of a restriction enzyme *KpnI* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction

at 37°C for 2 hours. After the digestion reaction, the solution was subjected to 0.8% (w/v) agarose gel electrophoresis to purify a DNA fragment of about 3.5 Kb.

A 4.0 µl portion of the obtained *EcoRV*-*KpnI* fragment (about 1.5 Kb) derived from the plasmid pKOFUT8gE2-1, 1.0 µl of the *HpaI*-*KpnI* fragment (about 3.5 Kb) derived from the plasmid ploxPPuro and 5.0 µl of Ligation High (manufactured by Toyobo) were mixed and allowed to undergo the ligation reaction at 16°C for 30 minutes. *E. coli* DH5α was transformed using the reaction solution, and a plasmid DNA was isolated in accordance with a known method from the obtained ampicillin-resistant clones. Herein, the plasmid is referred to pKOFUT8gE2-2.

(4) Construction of plasmid pscFUT8gE2-3

A plasmid pscFUT8gE2-3 was constructed by the following procedure (Fig. 36), using the plasmid pFUT8fgE2-4 obtained in Example 12(2) having a genome region comprising exon 2 of Chinese hamster FUT8.

In 35 µl of NEBuffer 1 (manufactured by New England Biolabs), 2.0 µg of the plasmid pFUT8fgE2-4 was dissolved, and 20 units of a restriction enzyme *HpaII* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. A DNA fragment was recovered from the reaction solution by ethanol precipitation, and then the DNA termini were changed to blunt ends using Blunting High (manufactured by Toyobo) in accordance with the manufacture's instructions. The DNA fragment was recovered by carrying out phenol/chloroform extraction and ethanol precipitation and dissolved in 35 µl of NEBuffer 2 (manufactured by New England Biolabs), and 20 units of a restriction enzyme *HindIII* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction,

the solution was subjected to 0.8% (w/v) agarose gel electrophoresis to purify a DNA fragment of about 3.5 Kb.

On the other hand, 1.0 µg of a plasmid LITMUS39 (manufactured by New England Biolabs) was dissolved in 35 µl of NEBuffer 2 (manufactured by New England Biolabs), and the solution was mixed with 20 units of a restriction enzyme *EcoRV* (manufactured by New England Biolabs) and 20 units of a restriction enzyme *HindIII* (manufactured by New England Biolabs) and subjected to the digestion reaction at 37°C for 2 hours. After the digestion reaction, the solution was subjected to 0.8% (w/v) agarose gel electrophoresis to purify a DNA fragment of about 2.8 Kb.

The obtained *HpaII-HindIII* fragment (4.0 µl, about 3.5 Kb) derived from the plasmid pFUT8fgE2-4, 1.0 µl of the *EcoRV-HindIII* fragment (about 2.8 Kb) derived from the plasmid LITMUS39 and 5.0 µl of Ligation High (manufactured by Toyobo) were mixed, followed by ligation reaction at 16°C for 30 minutes. *E. coli* DH5α was transformed using the reaction solution, and a plasmid DNA was isolated in accordance with a known method from the obtained ampicillin-resistant clones. Herein, the plasmid is referred to pscFUT8gE2-3.

(5) Construction of plasmid pKOFUT8gE2-3

A plasmid pKOFUT8gE2-3 was constructed by the following procedure (Fig. 37), using the plasmid pFUT8fgE2-4 obtained in Example 12(2) having a genome region comprising exon 2 of Chinese hamster FUT8.

In 35 µl of NEBuffer for *EcoRI* (manufactured by New England Biolabs), 2.0 µg of the plasmid pFUT8fgE2-4 was dissolved, and 20 units of a restriction enzyme *EcoRI* (manufactured by New England Biolabs) and 20 units of a restriction enzyme *HindIII* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, the

ethanol precipitation and dissolved in 30 μ l of NEBuffer 2 (manufactured by New England Biolabs) containing 100 μ g/ml of BSA (manufactured by New England Biolabs), and 20 units of a restriction enzyme *Hind*III (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, the solution was subjected to 0.8% (w/v) agarose gel electrophoresis to purify a DNA fragment of about 3.6 Kb.

Separately, 1.0 μ g of the plasmid pKOFUT8gE2-3 was dissolved in 35 μ l of NEBuffer for *Sal*I (manufactured by New England Biolabs), and 20 units of a restriction enzyme *Sal*I (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. A DNA fragment was recovered from the reaction solution by ethanol precipitation and dissolved in 35 μ l of NEBuffer 2 (manufactured by New England Biolabs), and 20 units of a restriction enzyme *Hind*III (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, 35 μ l of 1 mol/l Tris-HCl buffer (pH 8.0) and 3.5 μ l of *E. coli* Cl5-derived alkaline phosphatase (manufactured by Takara Shuzo) were added thereto, followed by reaction at 65°C for 30 minutes to dephosphorylate the DNA termini. After the dephosphorylation treatment, a DNA fragment was recovered by carrying out phenol/chloroform extraction and ethanol precipitation and dissolved in 10 μ l of sterile water.

The obtained *Sal*I-*Hind*III fragment (4.0 μ l, about 3.1 Kb) derived from the plasmid pscFUT8gE2-3, 1.0 μ l of the *Sal*I-*Hind*III fragment (about 4.8 Kb) derived from the plasmid pKOFUT8gE2-3 and 5.0 μ l of Ligation High (manufactured by Toyobo) were mixed, followed by ligation reaction at 16°C for 30 minutes. *E. coli* DH5 α was transformed using the reaction solution, and a plasmid DNA was isolated in accordance with a known method from the

obtained ampicillin-resistant clones. Herein, the plasmid is referred to pKOFUT8gE2-4.

(7) Construction of plasmid pKOFUT8gE2-5

A plasmid pKOFUT8gE2-5 was constructed by the following procedure (Fig. 39), using the plasmids pKOFUT8gE2-2 and pKOFUT8gE2-4 obtained in the items (3) and (6).

In 30 μ l of NEBuffer 4 (manufactured by New England Biolabs), 1.0 μ g of the plasmid pKOFUT8gE2-2 was dissolved, and 20 units of a restriction enzyme *Sma*I (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 25°C for 2 hours. A DNA fragment was recovered from the reaction solution by ethanol precipitation and dissolved in 30 μ l of NEBuffer 2 (manufactured by New England Biolabs), and 20 units of a restriction enzyme *Bam*HI (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, 30 μ l of 1 mol/l Tris-HCl buffer (pH 8.0) and 3.0 μ l of *E. coli* C15-derived alkaline phosphatase (manufactured by Takara Shuzo) were added thereto, followed by reaction at 65°C for 1 hour to dephosphorylate the DNA termini. After the dephosphorylation treatment, the DNA fragment was recovered by carrying out phenol/chloroform extraction and ethanol precipitation and dissolved in 10 μ l of sterile water.

Separately, 1.0 μ g of the plasmid pKOFUT8gE2-4 was dissolved in 30 μ l of NEBuffer 4 (manufactured by New England Biolabs), and 20 units of a restriction enzyme *Sma*I (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 25°C for 2 hours. A DNA fragment was recovered from the reaction solution by ethanol precipitation and dissolved in 30 μ l of NEBuffer 2 (manufactured by New England Biolabs), and 20 units of a restriction enzyme *Bam*HI (manufactured by New England

Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, the solution was subjected to 0.8% (w/v) agarose gel electrophoresis to purify a DNA fragment of about 5.2 Kb.

The obtained *Sma*I-*Bam*HI fragment (0.5 µl, about 5.0 Kb) derived from the plasmid pKOFUT8gE2-2, 4.5 µl of the *Sma*I-*Bam*HI fragment (about 5.4 Kb) derived from the plasmid pKOFUT8gE2-4 and 5.0 µl of Ligation High (manufactured by Toyobo) were mixed, followed by ligation reaction at 16°C for 15 hours. *E. coli* DH5α was transformed using the reaction solution, and a plasmid DNA was isolated in accordance with a known method from the obtained ampicillin-resistant clones. Herein, the plasmid is referred to pKOFUT8gE2-5.

(8) Construction of plasmid pKOFUT8Puro

A plasmid pKOFUT8Puro was constructed by the following procedure (Fig. 40), using the plasmid pKOFUT8gE2-5 obtained in the item (7).

In 50 µl of NEBuffer 4 (manufactured by New England Biolabs), 1.0 µg of a plasmid pKSelectDT (manufactured by Lexicon) was dissolved, and 16 units of a restriction enzyme *Rsr*II (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, the solution was subjected to 0.8% (w/v) agarose gel electrophoresis to purify a DNA fragment of about 1.2 Kb comprising a diphtheria toxin expression unit.

Separately, 1.0 µg of the plasmid pKOFUT8gE2-5 was dissolved in 50 µl of NEBuffer 4 (manufactured by New England Biolabs), and 16 units of a restriction enzyme *Rsr*II (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 2 hours. After the digestion reaction, 30 µl of 1 mol/l Tris-HCl buffer (pH 8.0) and 3.0 µl of *E. coli* C15-derived alkaline

phosphatase (manufactured by Takara Shuzo) were added thereto, followed by reaction at 65°C for 1 hour to dephosphorylate the DNA termini. After the dephosphorylation treatment, the DNA fragment was recovered by carrying out phenol/chloroform extraction and ethanol precipitation and dissolved in 10 µl of sterile water.

The obtained RsrII-RsrII fragment (1.0 µl, about 1.2 Kb) derived from the plasmid pKSelectDT, 1.0 µl of the RsrII-RsrII fragment (about 10.4 Kb) derived from the plasmid pKOFUT8gE2-5, 3.0 µl of sterile water and 5.0 µl of Ligation High (manufactured by Toyobo) were mixed, followed by ligation reaction at 16°C for 30 minutes. *E. coli* DH5α was transformed using the reaction solution, and a plasmid DNA was isolated in accordance with a known method from the obtained ampicillin-resistant clones. Herein, the plasmid is referred to pKOFUT8Puro.

2. Preparation of CHO cell in which one copy of the genome region containing α-1,6-fucosyltransferase (FUT8) gene exon 2 was disrupted

(1) Introduction of targeting vector

A Chinese hamster FUT8 genome region targeting vector pKOFUT8Puro constructed in the item 1 of this Example was introduced into the strain 5-03 prepared in the item 1(2) of Example 8.

A gene of the plasmid pKOFUT8Puro was introduced into the strain 5-03 as described below in accordance with the electroporation method [Cytotechnology, 3, 133 (1990)]. First, 150 µg of the plasmid pKOFUT8Puro was dissolved in 1.8 ml of NEBuffer for SalI (manufactured by New England Biolabs), and 600 units of a restriction enzyme SalI (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 5 hours to obtain a linear fragment. The reaction solution was extracted with phenol/chloroform extraction, followed by

ethanol precipitation, and the recovered linear plasmid was made into a 1 µg/µl aqueous solution. Separately, the strain 5-03 was suspended in a K-PBS buffer (137 mmol/l KCl, 2.7 mmol/l NaCl, 8.1 mmol/l Na₂HPO₄, 1.5 mmol/l KH₂PO₄, 4.0 mmol/l MgCl₂) to give a density of 8×10^7 cells/ml. After mixing 200 µl of the cell suspension (1.6×10^6 cells) with 4 µl (4 µg) of the linear plasmid, an entire volume of the cell-DNA mixture was transferred into Gene Pulser Cuvette (inter-electrode distance, 2 mm) (manufactured by BIO-RAD) and then the electroporation was carried out using a cell fusion apparatus Gene Pulser (manufactured by BIO-RAD) at 350 V pulse voltage and 250 µF electric capacity. After carrying out the electroporation using 30 cuvettes in the same manner, the cell suspension was suspended in IMDM medium (manufactured by Life Technologies) supplemented with 10% fetal bovine serum (manufactured by Life Technologies) and 1 × concentration HT supplement (manufactured by Life Technologies) and inoculated into 30 adhesion cell culture dishes of 10 cm in diameter (manufactured by Falcon). After culturing them at 37°C for 24 hours in 5% CO₂, the culture supernatant was removed, and IMDM medium (manufactured by Life Technologies) supplemented with 15 µg/ml puromycin (manufactured by SIGMA) and 10% fetal bovine dialyzed serum (manufactured by Life Technologies) was dispensed in 10 ml portions. After culturing them for 10 days while repeating the medium exchange at intervals of 3 to 4 days, puromycin-resistant cell lines were obtained.

(2) Preparation of targeting vector-introduced cell lines

Arbitrary 900 colonies were obtained as follows from the puromycin-resistant cell lines obtained in the item (1).

First, culture supernatant was removed from the 10 cm dish in which colony of puromycin-resistant cell lines

(manufactured by SIGMA) and 10% fetal bovine dialyzed serum (manufactured by Life Technologies), the cells were inoculated into a 6 well flat bottom plate for adhesion cell (manufactured by Greiner). Genome DNA was prepared from the clone in the plate in accordance with a known method [Nucleic Acids Research, 3, 2303 (1976)] and dissolved overnight in 150 μ l of a TE-RNase buffer (pH 8.0) (10 mmol/l Tris-HCl, 1 mmol/l EDTA, 200 μ g/ml RNase A).

In 120 μ l of NEBuffer 3 (manufactured by New England Biolabs), 12 μ g of the obtained genome DNA was dissolved, and 25 units of a restriction enzyme *Pst*I (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C overnight. A DNA fragment was recovered from the reaction solution by ethanol precipitation, dissolved in 20 μ l of TE buffer (pH 8.0) (10 mmol/l Tris-HCl, 1 mmol/l EDTA) and then subjected to 0.8% (w/v) agarose gel electrophoresis. After the electrophoresis, the genome DNA was transferred onto a nylon membrane in accordance with a known method [Proc. Natl. Acad. Sci. USA, 76, 3683 (1979)]. After completion of the transfer, the nylon membrane was heated at 80°C for 2 hours.

Separately, a probe used in the Southern blotting was prepared as follows. First, primers (SEQ ID NOs:28 and 29) which bind to a sequence outside the targeting vector homologous region with the FUT8 genome region obtained in Example 12 were designed. Next, using a DNA polymerase ExTaq (manufactured by Takara Shuzo), 20 μ l of a reaction solution [ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs and 0.5 μ mol/l gene-specific primers (SEQ ID NOs:28 and 29)] containing 4.0 ng of the plasmid pFUT8fgE2-2 obtained in Example 12(2) was prepared, and polymerase chain reaction (PCR) was carried out. The PCR was carried out by heating at 94°C for 1 minute and subsequent 25 cycles of heating at 94°C for 30 seconds, 55°C for 30

twice, the membrane was soaked in 50 ml of 0.2 x SSC-0.1% (w/v) SDS and heated at 65°C for 15 minutes. After washing the nylon membrane, it was exposed to an X-ray film two nights at -80°C for development.

By the restriction enzyme *Bam*HI treatment, a DNA fragment of about 19.0 Kb was formed from a wild type FUT8 allele. Also, a DNA fragment of about 12.5 Kb was formed from an allele in which homologous recombination with a targeting vector was generated. In addition, when the puromycin resistance gene (about 1.5 Kb) was deleted from the allele in which homologous recombination was generated, a DNA fragment of about 11.0 Kb was formed by the same treatment.

By the method, such specific fragments of about 19.0 Kb and about 11.0 Kb were found from the genome DNA of 5 clones among the 6 clones. Since the quantitative ratio of both fragments was 1 : 1, it was shown that the puromycin resistance gene was deleted from the cell lines in which 1 copy of the FUT8 genome region was disrupted. Hereinafter, one of the clone is called 1st.ΔFUT8 2-46-1. Also, results of the genome Southern blotting of the strain 1st.ΔFUT8 2-46-1, 1st.ΔFUT8 2-46 and 5-03 are shown in Fig. 41. Also, the strain 1st.ΔFUT8 2-46-1, as a name of 2-46-1, has been deposited on September 26, 2001, as FERM BP-7755 in International Patent Organism Depositary, National Institute of Advanced Industrial Science and Technology (AIST Tsukuba Central 6, 1-1, Higashi 1-Chome Tsukuba-shi, Ibaraki-ken 305-8566 Japan)).

4. Purification of antibody produced by α-1,6-fucosyltransferase (FUT8) gene-disrupted cell line

1st.ΔFUT8 2-46-1 obtained in the item 3 of this Example by disrupting one copy of the FUT8 allele was suspended in IMDM medium (manufactured by Life Technologies) supplemented with 15 μg/ml of puromycin

8535 8535 0612

centrifugation tubes, combined and then suspended in 30 ml of RPMI1640-FBS(10). After centrifugation at room temperature at 1,200 rpm for 15 minutes, the supernatant was discarded and the cells were suspended in 20 ml of RPMI1640-FBS(10). The washing step was repeated twice and then 2.5×10^6 cells/ml of an effector cell suspension was prepared using RPMI1640-FBS(10).

The target cell suspension was dispensed at 50 μ l (1×10^4 cells/well) into each well of a 96 well U-bottom plate (manufactured by Falcon). Subsequently, the effector cell suspension was dispensed at 100 μ l (2.5×10^5 cells/well) into each well to thereby adjust the ratio of the effector cells to the target cells to 25 : 1. Next, using RPMI1640-FBS(10), a series of dilution solution of 0.01 μ g/ml, 0.1 μ g/ml, 1 μ g/ml and 10 μ g/ml was prepared from each of the anti-CCR4 antibodies obtained in the item 5 of Example 13, and the diluted solutions were dispensed in 50 μ l portions into the wells to give final concentrations of 0.0025 μ g/ml, 0.025 μ g/ml, 0.25 μ g/ml and 2.5 μ g/ml, respectively. After the reaction at 37°C for 4 hours in 5% CO₂, the plate was centrifuged at 1,200 rpm for 5 minutes. Into a 12 mm diameter RIA tube (manufactured by IWAKI), 75 μ l of the supernatant in each well was batched off and the amount of the dissociated ⁵¹Cr was measured using MINAX- γ auto-gamma counter 5550 (manufactured by PACKARD).

Also, the amount of the spontaneously dissociated ^{51}Cr was calculated by carrying out the same reaction in a reaction mixture in which 150 μl of RPMI1640-FBS(10) was added instead of the effector cell suspension and antibody solution. The amount of the total dissociated ^{51}Cr was calculated by carrying out the same reaction in a reaction mixture in which 100 μl of 1 N hydrochloric acid and 50 μl of RPMI1640-FBS(10) were added instead of the effector cell

number of cells to be treated. Hereinafter, these cell lines were used in analyses.

(2) Preparation of anti-CCR4 human chimeric antibody-producing cell

An anti-CCR4 human chimeric antibody expression plasmid pKANTEX2160 was introduced into the three lectin-resistant cell lines obtained in the (1) by the method described in Example 8, and gene amplification by a drug MTX was carried out to prepare an anti-CCR4 human chimeric antibody-producing cell line. By measuring an amount of antibody expression by the ELISA described in Example 8-2, antibody-expressing transformants were obtained from each of the CHO-LCA, CHO-AAL and CHO-PHA. Regarding each of the obtained transformants, a transformant derived from CHO-LCA was named CHO/CCR4-LCA, a transformant derived from CHO-AAL was named CHO/CCR4-AAL and a transformant derived from CHO-PHA was named CHO/CCR4-PHA. Also, the CHO/CCR4-LCA, as a name of Nega-13, has been deposited on September 26, 2001, as FERM BP-7756 in International Patent Organism Depository, National Institute of Advanced Industrial Science and Technology (AIST Tsukuba Central 6, 1-1, Higashi 1-Chome Tsukuba-shi, Ibaraki-ken 305-8566 Japan)).

(3) Production of potent ADCC activity antibody by lectin-resistant CHO cell

Using the three transformants obtained in the (2), purified antibodies were obtained by the method described in Example 8-3. The antigen binding activity of each of the purified anti-CCR4 human chimeric antibodies was evaluated using the ELISA described in Example 8-2. The antibodies produced by all transformants showed an antigen binding activity similar to that of the antibody produced by a recombinant cell line (strain 5-03) prepared in Example 8 using general CHO/DG44 cell as the host. Using

these purified antibodies, ADCC activity of each of the purified anti-CCR4 human chimeric antibodies was evaluated in accordance with the method described in Example 8-7. The results are shown in Fig. 43. In comparison with the antibody produced by the strain 5-03, about 100 folds-increased ADCC activity was observed in the antibodies produced by the CHO/CCR4-LCA and CHO/CCR4-AAL. On the other hand, no significant increase in the ADCC activity was observed in the antibody produced by the CHO/CCR4-PHA. Also, when ADCC activities of the antibodies produced by the CHO/CCR4-LCA and YB2/0 were compared in accordance with the method described in Example 8-7, it was found that the antibody produced by the CHO/CCR4-LCA shows more potent ADCC activity than the antibody produced by the strain 5-03, similar to the case of the antibody KM2760-1 produced by the YB2/0 cell line prepared in Example 8-1 (Fig. 44).

(4) Sugar chain analysis of antibodies produced by lectin-resistant CHO cell

Sugar chains of the anti-CCR4 human chimeric antibodies purified in the (3) were analyzed. The solution of each of the purified antibodies was exchanged to 10 mM KH_2PO_4 using Ultra Free 0.5-10K (manufactured by Millipore). The exchange was carried out in such a manner that the exchanging ratio became 80 folds or more. The concentration of the antibodies after the solution exchange was measured using UV-1600 (manufactured by Shimadzu). The molar absorption coefficient was calculated from the amino acid sequence of each antibody based on the following equation (III) [Advances in Protein Chemistry, 12, 303 (1962)], and the concentration was determined by defining the absorbance at 280 nm as 1.38 mg/ml.

$$E_{1\text{mol/l}} = A \times n1 + B \times n2 + C \times n3 \quad (\text{III})$$

$$E_{1\text{mol/ml}} = E_{1\text{mol/l}}/\text{MW}$$

frozen at -80°C in a freezer and freeze-dried for about 17 hours. Sugar chains were recovered from the freeze-dried sample using Cellulose Cartridge Glycan Preparation Kit manufactured by Takara Syuzo Co., Ltd. The sample sugar chain solution was dried using a centrifugal evaporator and then subjected to fluorescence labeling with 2-aminopyridine [*J. Biochem.*, 95, 197 (1984)]. The 2-aminopyridine solution was prepared by adding 760 μl of HCl per 1 g of 2-aminopyridine (1 \times PA solution) and diluting the solution 10 folds with reverse osmosis purified water (10-folds diluted PA solution). The sodium cyanoborohydride solution was prepared by adding 20 μl of 1 \times PA solution and 430 μl of reverse osmosis purified water per 10 mg of sodium cyanoborohydride. To the sample, 67 μl of a 10 folds-diluted PA solution was added, followed by reaction at 100°C for 15 minutes and spontaneously cooled, and 2 μl of sodium cyanoborohydride was further added thereto, followed by reaction at 90°C for 12 hours for fluorescence labeling of the sample sugar chains. The fluorescence-labeled sugar chain group (PA-treated sugar chain group) was separated from excess reagent using Superdex Peptide HR 10/30 column (manufactured by Pharmacia). This step was carried out using 10 mM ammonium bicarbonate as the eluent at a flow rate of 0.5 ml/min and at a column temperature of room temperature, and using a fluorescence detector of 320 nm excitation wavelength and 400 nm fluorescence wavelength. The eluate was recovered 20 to 30 minutes after addition of the sample and dried using a centrifugal evaporator to be used as purified PA-treated sugar chains. Next, reverse phase HPLC analysis of the purified PA-treated sugar chains was carried out using CLC-ODS column (manufactured by Shimadzu, ϕ 6.0 mm \times 159 mm). The step was carried out at a column temperature of 55°C and at a flow rate of 1 ml/min and using a fluorescence detector of 320 nm excitation wavelength and

400 nm fluorescence wavelength. The column was equilibrated with a 10 mM sodium phosphate buffer (pH 3.8) and elution was carried out for 80 minutes by a 0.5% 1-butanol linear density gradient. Each of the PA-treated sugar chain was identified by post source decay analysis of each peak of the separated PA-treated sugar chains using matrix-assisted laser ionization time of flight mass spectrometry (MALDI-TOF-MS analysis), comparison of elution positions with standards of PA-treated sugar chain manufactured by Takara Syuzo, and reverse phase HPLC analysis after digestion of each PA-treated sugar chain using various enzymes (Fig. 45). Each of the sugar chain content was calculated from each of the peak area of PA-treated sugar chain by reverse HPLC analysis. A PA-treated sugar chain whose reducing end is not *N*-acetylglucosamine was excluded from the peak area calculation, because it is an impurity or a by-product during preparation of PA-treated sugar chain.

The analysis was carried out in the same manner as in Example 11(6) using a sodium phosphate buffer (pH 3.8) as buffer A and a sodium phosphate buffer (pH 3.8) + 0.5% 1-butanol as buffer B.

In Fig. 45, the ratio of the α -1,6-fucose-free sugar chain group was calculated from the area occupied by the peaks (i) to (iv) among (i) to (viii), and the ratio of the α -1,6-fucose-bound sugar chain group from the area occupied by the peaks (v) to (viii) among (i) to (viii).

Results of the sugar chain structure analysis of the purified anti-CCR4 human chimeric antibodies produced by lectin-resistant cell lines are shown in Table 6. The result shows the analysis of sugar chains of the anti-CCR4 human chimeric antibody produced by lectin-resistant cell lines. The ratio of α -1,6-fucose-free sugar chains (%) calculated from peak areas by analyzing by the method described in Example d(4) is shown in the table.

Table 6

Antibody producing cells	α -1,6-Fucose-free complex double-chain sugar chain (%)
Strain 5-03	9
Strain CHO/CCR4-LCA	48
Strain CHO/CCR4-AAL	27
Strain CHO/CCR4-PHA	8

In comparison with the antibody produced by the strain 5-03, the ratio of the α -1,6-fucose-free sugar chains was increased from 9% to 48% in the antibody produced by the CHO/CCR4-LCA when calculated from the analyzed peak area. The ratio of α -1,6-fucose-free sugar chains was increased from 9% to 27% in the antibody produced by the CHO/CCR4-AAL. On the other hand, changes in the sugar chain pattern and ratio of the α -1,6-fucose-free sugar chains were hardly found in the PHA-resistant cell line when compared with the strain 5-03.

Example 15

Analysis of lectin-resistant CHO cell line:

1. Analysis of expression level of GMD enzyme in an anti-CCR4 human chimeric antibody-producing cell line CHO/CCR4-LCA

The expression level of each of the genes of GMD (GDP-mannose 4,6-dehydratase), GFPP (GDP-keto-6-deoxymannose 3,5-epimerase, 4-reductase) and FX (GDP-beta-L-fucose pyrophosphorylase) known as fucose biosynthesis enzymes and FUT8 (α -1,6-fucosyltransferase) as a fucose transferase, in the anti-CCR4 human chimeric antibody-producing cell line CHO/CCR4-LCA obtained in Example 14, was analyzed using RT-PCR method.

(1) Preparation of RNA from various cell lines

Each of the CHO/DG44 cell, the anti-CCR4 human chimeric antibody-producing cell line 5-03 obtained in Example 8-1(2) and the anti-CCR4 human chimeric antibody-producing cell line CHO/CCR4-LCA obtained in Example 14(2) was subcultured at 37°C in a 5% CO₂ incubator and then cultured for 4 days. After culturing them, total RNA was prepared from 1×10^7 cells of each cell line using RNeasy Protect Mini Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions. Subsequently, single-stranded cDNA was synthesized from 5 µg of each RNA in a 20 µl of a reaction solution using SUPER SCRIPT First-Strand Synthesis System for RT-PCR (manufactured by GIBCO BRL) in accordance with the manufacture's instructions.

(2) Analysis of expression level of GMD gene using RT-PCR

In order to amplify GMD cDNA by PCR, a 24 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:32 and a 26 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:33 were prepared based on the CHO cell-derived GMD cDNA sequence shown in Example 17-1.

Next, 20 µl of a reaction solution [1 × Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mM dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 µM of the synthetic DNA primers of SEQ ID NOs:32 and 33] containing 0.5 µl of the single-stranded cDNA prepared from each cell line in the item (1) as the template was prepared, and PCR was carried out using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 30 cycles of heating of 94°C for 1 minute and 68°C for 2 minutes as one cycle. After subjecting 10 µl of the PCR reaction solution to agarose electrophoresis, DNA fragments were stained using Cyber Green (manufactured by BMA) and then the amount

of the DNA fragment of about 350 bp was measured using Fluor Imager SI (manufactured by Molecular Dynamics).

(3) Analysis of expression level of GFPP gene using RT-PCR

In order to amplify GFPP cDNA by PCR, a 27 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:34 and a 23 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:35 were prepared based on the CHO cell-derived GFPP cDNA sequence obtained in Example 16-2.

Next, 20 μ l of a reaction solution [1 \times Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mM dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 μ M of the synthetic DNA primers of SEQ ID NOs:34 and 35] containing 0.5 μ l of the single-stranded cDNA prepared from each cell line in the item (1) as the template was prepared, and PCR was carried out using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 24 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle. After subjecting 10 μ l of the PCR reaction solution to agarose electrophoresis, DNA fragments were stained using Cyber Green (manufactured by BMA) and then the amount of the DNA fragment of about 600 bp was measured using Fluor Imager SI (manufactured by Molecular Dynamics).

(4) Analysis of expression level of FX gene using RT-PCR

In order to amplify FX cDNA by PCR, a 28 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:36 and a 28 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:37 were prepared based on the CHO cell-derived FX cDNA sequence shown in Example 16-1.

Next, 20 μ l of a reaction solution [1 \times Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mM dNTPs, 0.5

Table 7

	GMD	GEPP	FX	FUT8
Strain CHO/DG44	1	1	1	1
Strain 5-03	1.107	0.793	1.093	0.901
Strain 5-03-derived LCA-resistant cell CHO/CCR4-LCA	0.160	0.886	0.920	0.875

As shown in Table 7, the expression level of GMD gene in the CHO/CCR4-LCA was decreased to about 1/10 in comparison with other cell lines. In this case, the test was independently carried out twice, and the average value was used.

2. Analysis using anti-CCR4 human chimeric antibody-producing CHO/CCR4-LCA in which GMD gene was forced to express

(1) Construction of CHO cell-derived GMD gene expression vector pAGE249GMD

Based on the CHO cell-derived GMD cDNA sequence obtained in Example 17-1, a 28 mer primer having the nucleotide sequence shown by SEQ ID NO:38 and a 29 mer primer having the nucleotide sequence shown by SEQ ID NO:39 were prepared. Next, 20 µl of a reaction solution [1 × Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mM dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 µM of the synthetic DNA primers of SEQ ID NOs:38 and 39] containing 0.5 µl of the CHO cell-derived GMD single-stranded cDNA prepared in the item 1(1) of this Example as the template was prepared, and PCR was carried out using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequently 8 cycles of heating at 94°C for 1 minute, 58°C for 1 minute and 72°C for 1 minute as one cycle, and then 22 cycles of heating at 94°C for 1 minute and 68°C as one cycle. After completion of the reaction, the PCR reaction solution was

fractionated by agarose electrophoresis, and then a DNA fragment of about 600 bp was recovered using Gene Clean II Kit (manufactured by BIO 101) in accordance with the manufacture's instructions. The recovered DNA fragment was connected to pT7Blue(R) vector (manufactured by Novagen) using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5 α (manufactured by Toyobo) was transformed using the obtained recombinant plasmid DNA to obtain a plasmid mt-C (cf. Fig. 46).

Next, based on the CHO cell-derived GMD cDNA sequence obtained in Example 17-1, a 45 mer primer having the nucleotide sequence shown by SEQ ID NO:40 and a 31 mer primer having the nucleotide sequence shown by SEQ ID NO:41 were prepared. Next, 20 μ l of a reaction solution [1 \times Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mM dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 μ M of the synthetic DNA primers of SEQ ID NOs:40 and 41] containing 0.5 μ l of the CHO cell-derived GMD single-stranded cDNA prepared in the item 1(1) of this Example as the template was prepared, and PCR was carried out using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequently 8 cycles of heating at 94°C for 1 minute, 57°C for 1 minute and 72°C for 1 minute as one cycle, and then 22 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle. After completion of the reaction, the PCR reaction solution was fractionated by agarose electrophoresis, and then a DNA fragment of about 150 bp was recovered using Gene Clean II Kit (manufactured by BIO 101) in accordance with the manufacture's instructions. The recovered DNA fragment was connected to pT7Blue(R) vector (manufactured by Novagen) using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5 α (manufactured by Toyobo) was transformed using the obtained recombinant plasmid DNA to obtain a plasmid ATG (cf. Fig. 47).

Next, 2 µg of the plasmid WT-N(-) was allowed to react with a restriction enzyme *Bam*HI (manufactured by Takara Shuzo) at 37°C for 16 hours, DNA was recovered by carrying out phenol/chloroform extraction and ethanol precipitation and allowed to react with a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) at 37°C for 16 hours, the digest was fractionated by agarose electrophoresis and then a DNA fragment of about 1 kbp was recovered using Gene Clean II Kit (manufactured by BIO 101) in accordance with the manufacture's instructions. The plasmid pBluescript SK(-) (3 µg; manufactured by

for 19 days to obtain colonies of hygromycin-resistant transformants.

In the same manner, the pAGE249 vector was introduced into the CHO/CCR4-LCA by the same method to obtain colonies of hygromycin-resistant transformants.

(3) Culturing of GMD gene-expressed CHO/CCR4-LCA and purification of antibody

Using IMDM-dFBS(10) medium comprising 200 nM MTX (manufactured by SIGMA) and 0.5 mg/ml hygromycin, the GMD-expressing transformant cells obtained in the item (2) were cultured using a 182 cm² flask (manufactured by Greiner) at 37°C in a 5% CO₂ incubator. Several days thereafter, when the cell density reached confluent, the culture supernatant was discarded and the cells were washed with 25 ml of PBS buffer (manufactured by GIBCO BRL) and mixed with 35 ml of EXCELL301 medium (manufactured by JRH). After culturing them at 37°C in a 5% CO₂ incubator for 7 days, the culture supernatant was recovered. An anti-CCR4 chimeric antibody was purified from the culture supernatant using Prosep-A (manufactured by Millipore) in accordance with the manufacture's instructions.

In the same manner, the pAGE249 vector-introduced transformant cells were cultured by the same method and then anti-CCR4 chimeric antibody was recovered and purified from the culture supernatant.

(4) Measurement of lectin resistance in transformed cells

The GMD-expressing transformant cells obtained in the item (2) were suspended in IMDM-dFBS(10) medium comprising 200 nM MTX (manufactured by SIGMA) and 0.5 mg/ml hygromycin to give a density of 6×10^4 cells/ml, and the suspension was dispensed in 50 µl/well portions into a 96 well culture plate (manufactured by Iwaki Glass). Next, a medium prepared by suspending at concentrations of 0 mg/ml,

(5) *In vitro* cytotoxic activity (ADCC activity) of anti-CCR4 chimeric antibody obtained from GMD-expressed CHO/CCR4-LCA

In order to evaluate *in vitro* cytotoxic activity of the purified anti-CCR4 chimeric antibody obtained in the item (3), the ADCC activity was measured in accordance with the following methods.

i) Preparation of target cell suspension

A 3.7 MBq equivalent of a radioactive substance $\text{Na}_2^{51}\text{CrO}_4$ was added to 1×10^6 cells of the CCR4-EL4 (cf. Example 8-7) cultured using a medium prepared by adding 500 $\mu\text{g}/\text{ml}$ G418 sulfate (manufactured by Nakalai Tesque) to the RPMI1640-FBS(10) medium, followed by reaction at 37°C for 90 minutes to thereby label the cells with a radioisotope. After the reaction, the cells were washed three times by suspension in the RPMI1640-FBS(10) medium and subsequent centrifugation, re-suspended in the medium and then incubated at 4°C for 30 minutes in ice for spontaneous dissociation of the radioactive substance. After centrifugation, the cells were adjusted to 2.5×10^5 cells/ml by adding 5 ml of the RPMI1640-FBS(10) medium and used as a target cell suspension.

ii) Preparation of effector cell suspension

From a healthy person, 50 ml of venous blood was collected and gently mixed with 0.5 ml of heparin sodium (manufactured by Takeda Pharmaceutical). Using Lymphoprep (manufactured by Nycomed Pharma AS), the mixture was centrifuged in accordance with the manufacture's instructions to separate a mononuclear cell layer. The cells were washed three times by centrifuging using the RPMI1640-FBS(10) medium and then resuspended in the medium to give a density of 2×10^6 cells/ml and used as a effector cell suspension.

in the CHO/CCR4-LCA is decreased and, as a result, an antibody having potent ADCC activity can be produced

(6) Sugar chain analysis of anti-CCR4 chimeric antibody derived from GMD-expressed CHO/CCR4-LCA

Sugar chains binding to the purified anti-CCR4 chimeric antibody obtained in the item (3) were analyzed in accordance with the method shown in Example 14(4), with the analyzed results shown in Fig. 55. In comparison with the purified anti-CCR4 chimeric antibody prepared from CHO/CCR4-LCA in Example 14, the ratio of sugar chain having no α -1,6-fucose in the purified anti-CCR4 chimeric antibody derived from GMD-expressed CHO/CCR4-LCA was decreased to 9% when calculated from the peak area. Thus, it was shown that the ratio of sugar chain having no α -1,6-fucose in the antibody produced by the cell is decreased to similar level of the antibody produced by the strain 5-03, by expressing GMD gene in the CHO/CCR4-LCA

Example 16

Preparation of various genes encoding enzymes relating to the sugar chain synthesis in CHO cell:

1. Determination of CHO cell-derived FX cDNA sequence

(1) Extraction of total RNA from CHO/DG44 cell

CHO/DG44 cells were suspended in IMDM medium containing 10% fetal bovine serum (manufactured by Life Technologies) and 1 x concentration HT supplement (manufactured by Life Technologies), and 15 ml of the suspension was inoculated into a T75 flask for adhesion cell culture use (manufactured by Greiner) to give a density of 2×10^5 cells/ml. On the second day after culturing them at 37°C in a 5% CO₂ incubator, 1×10^7 of the cells were recovered and total RNA was extracted therefrom using RNeasy (manufactured by QIAGEN) in accordance with the manufacture's instructions.

determined encodes open reading frame (ORF) partial sequence of the Chinese hamster FX.

(4) Synthesis of single-stranded cDNA for RACE

Single-stranded cDNA samples for 5' and 3' RACE were prepared from the CHO/DG44 total RNA extracted in the item (1) using SMARTTM RACE cDNA Amplification Kit (manufactured by CLONTECH) in accordance with the manufacture's instructions. In the case, PowerScriptTM Reverse Transcriptase (manufactured by CLONTECH) was used as the reverse transcriptase. Each single-stranded cDNA after the preparation was diluted 10 folds with the Tricin-EDTA buffer attached to the kit and used as the template of PCR.

(5) Determination of Chinese hamster-derived FX full length cDNA by RACE method

Based on the FX partial sequence derived from Chinese hamster determined in the item (3), primers FXGSP1-1 (SEQ ID NO:44) and FXGSP1-2 (SEQ ID NO:45) for the Chinese hamster FX-specific 5' RACE and primers FXGSP2-1 (SEQ ID NO:46) and FXGSP2-2 (SEQ ID NO:47) for the Chinese hamster FX-specific 3' RACE were designed.

Next, polymerase chain reaction (PCR) was carried out using Advantage2 PCR Kit (manufactured by CLONTECH), by preparing 50 µl of a reaction solution [Advantage2 PCR buffer (manufactured by CLONTECH), 0.2 mM dNTPs, 0.2 µmol/l Chinese hamster FX-specific primers for RACE and 1 × concentration of common primers (manufactured by CLONTECH)] containing 1 µl of the CHO/DG44-derived single-stranded cDNA for RACE prepared in the item (4).

The PCR was carried out by repeating 20 cycles of heating at 94°C for 5 seconds, 68°C for 10 seconds and 72°C for 2 minutes as one cycle.

subsequent 30 cycles of heating at 94°C for 1 minute, 58°C for 2 minutes and 72°C for 3 minutes as one cycle, and final heating at 72°C for 10 minutes.

After the PCR, the reaction solution was subjected to 2% agarose gel electrophoresis, and a specific amplified fragment of 1.4 Kbp was purified using QuiaexII Gel Extraction Kit (manufactured by Quiagen) and eluted with 20 µl of sterile water. Into a plasmid pCR2.1, 4 µl of the amplified fragment was employed to insert in accordance with the instructions attached to TOPO TA Cloning Kit (manufactured by Invitrogen), and E. coli DH5α was transformed using the reaction solution.

Plasmid DNAs were isolated from the appeared several kanamycin-resistant colonies to obtain 3 clones into which GFPP cDNA partial fragments were respectively integrated. They are referred to as GFPP clone 8, GFPP clone 11 and GFPP clone 12.

The nucleotide sequence of the cDNA inserted into each of the GFPP clone 8, GFPP clone 11 and GFPP clone 12 was determined using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction kit (manufactured by Parkin Elmer) in accordance with the method described in the manufacture's instructions. It was confirmed that the inserted cDNA whose sequence was determined encodes open reading frame (ORF) partial sequence of the Chinese hamster GFPP.

(2) Determination of Chinese hamster GFPP full length cDNA by RACE method

Based on the Chinese hamster FX partial sequence determined in the item 2(1), primers GFPP GSP1-1 (SEQ ID NO:52) and GFPP GSP1-2 (SEQ ID NO:53) for the Chinese hamster FX-specific 5' RACE and primers GFPP GSP2-1 (SEQ ID NO:54) and GFPP GSP2-2 (SEQ ID NO:55) for the Chinese hamster GFPP-specific 3' RACE were designed.

The nucleotide sequence of the cDNA of each of the clones obtained by the 5' and 3' RACE was determined using DNA Sequencer 377 (manufactured by Parkin Elmer) in accordance with the method described in the manufacture's instructions. By comparing the cDNA nucleotide sequences after the nucleotide sequence determination, reading errors of bases due to PCR were excluded and the full length nucleotide sequence of Chinese hamster GFPP cDNA was determined. The determined sequence is shown in SEQ ID NO:51.

Example 17

Preparation of CHO cell-derived GMD gene:

1. Determination of CHO cell-derived GMD cDNA sequence

(1) Preparation of CHO cell-derived GMD gene cDNA (preparation of partial cDNA excluding 5'- and 3'-terminal sequences)

Rodents-derived GMD cDNA was searched in a public data base (BLAST) using a human-derived GMD cDNA sequence (GenBank Accession No. AF042377) registered at GenBank as a query, and three kinds of mouse EST sequences were obtained (GenBank Accession Nos. BE986856, BF158988 and BE284785).

restriction enzyme *EcoRI* at 37°C for 16 hours, the digest was subjected to agarose electrophoresis and then a DNA fragment of about 150 bp was recovered using Gene Clean II Kit (manufactured by BIO 101) in accordance with the manufacture's instructions. The recovered DNA fragments were respectively subjected to terminal dephosphorylation using Calf Intestine Alkaline Phosphatase (manufactured by Takara Shuzo) and then ligated using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5α (manufactured by Toyobo) was transformed using the obtained recombinant plasmid DNA to obtain a plasmid CHO-GMD (cf. Fig. 54).

(2) Determination of 5'-terminal sequence of CHO cell-derived GMD cDNA

A 24 mer primer having the nucleotide sequence represented by SEQ ID NO:61 was prepared from 5'-terminal side non-coding region nucleotide sequences of CHO cell-derived human and mouse GMD cDNA, and a 32 mer primer having the nucleotide sequence represented by SEQ ID NO:62 from CHO cell-derived GMD cDNA sequence were prepared, and PCR was carried out by the following method to amplify cDNA. Then, 20 µl of a reaction solution [1 × Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mM dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 µM of the synthetic DNA primers of SEQ ID NO:61 and SEQ ID NO:62] containing 0.5 µl of the single-stranded cDNA prepared in Example 15-1(1) was prepared as the template, and the reaction was carried out therein using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes, subsequent 20 cycles of heating at 94°C for 1 minute, 55°C for 1 minute and 72°C for 2 minutes as one cycle and further 18 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle. After fractionation of the PCR reaction solution by agarose

μl of the cDNA for 3' RACE as the template was prepared, and PCR was carried out using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 30 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle.

After completion of the reaction, 1 μl of the PCR reaction solution was diluted 20 folds with Tricin-EDTA buffer (manufactured by CLONTECH). Then, 20 μl of a reaction solution [ExTaq buffer (manufactured by Takara Shuzo), 0.2 mM dNTPs, 0.5 unit of EX Taq polymerase (manufactured by Takara Shuzo), 0.5 μM of the 25 mer synthetic DNA primer shown in SEQ ID NO:64 [generated on the base of the CHO cell-derived GMD cDNA sequence determined in the item (1)] and 0.5 μM of Nested Universal Primer (attached to SMART™ RACE cDNA Amplification Kit; manufactured by CLONTECH) containing 1 μl of the 20 folds-diluted aqueous solution as the template] was prepared, and the reaction was carried out using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 30 cycles at 94°C for 1 minute and 68°C for 2 minutes as one cycle.

After completion of the reaction, the PCR reaction solution was fractionated by agarose electrophoresis and then a DNA fragment of about 700 bp was recovered using Gene Clean II Kit (manufactured by BIO 101) in accordance with the manufacture's instructions. The recovered DNA fragment was ligated to a pT7Blue(R) vector (manufactured by Novagen) using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5α (manufactured by Toyobo) was transformed using the obtained recombinant plasmid DNA, thereby obtaining a plasmid 3'GMD. Using DNA Sequencer 377 (manufactured by Parkin Elmer), the nucleotide sequence of 27 bases in upstream of the termination codon of CHO cell-derived GMD cDNA contained in the plasmid was determined.

